

Quantitative imaging of mitochondrial and cytosolic free zinc levels in an in vitro model of ischemia/reperfusion

Bryan J. McCranor · Rebecca A. Bozym ·
Michele I. Vitolo · Carol A. Fierke · Linda Bambrick ·
Brian M. Polster · Gary Fiskum · Richard B. Thompson

Received: 9 October 2011 / Accepted: 17 December 2011 / Published online: 20 March 2012
© Springer Science+Business Media, LLC 2012

Abstract The role of zinc ion in cytotoxicity following ischemic stroke, prolonged status epilepticus, and traumatic brain injury remains controversial, but likely is the result of mitochondrial dysfunction. We describe an excitation ratio-metric fluorescence biosensor based on human carbonic anhydrase II variants expressed in the mitochondrial matrix, permitting free zinc levels to be quantitatively imaged therein. We observed an average mitochondrial matrix free zinc concentration of 0.2pM in the PC12 rat pheochromocytoma cell culture line. Cytoplasmic and mitochondrial free zinc levels were imaged in a cellular oxygen glucose deprivation (OGD) model of ischemia/reperfusion. We observed a significant increase in mitochondrial zinc 1 h following 3 h

OGD, at a time point when cytosolic zinc levels were depressed. Following the increase, mitochondrial zinc levels returned to physiological levels, while cytosolic zinc increased gradually over a 24 h time period in viable cells. The increase in intramitochondrial zinc observed during reoxygenation after OGD may contribute to bioenergetic dysfunction and cell death that occurs with both in vitro and in vivo models of reperfusion.

Keywords Zinc · Oxidative stress · Mitochondria · Ischemia · Stroke · Fluorescence biosensor · Ratiometric · Carbonic anhydrase II

B. J. McCranor · R. A. Bozym · M. I. Vitolo ·
R. B. Thompson (✉)
Department of Biochemistry and Molecular Biology,
University of Maryland School of Medicine,
108 N. Greene St.,
Baltimore, MD 21201, USA
e-mail: rthompo@umaryland.edu

M. I. Vitolo
University of Maryland Marlene and Stewart Greenebaum NIH
Cancer Center, University of Maryland,
22 S. Greene St.,
Baltimore, MD 21201, USA

C. A. Fierke
Department of Chemistry, University of Michigan,
930 N. University Ave.,
Ann Arbor, MI 48109, USA

L. Bambrick · B. M. Polster · G. Fiskum
Department of Anesthesiology and the Center for Shock, Trauma,
and Anesthesiology Research,
University of Maryland School of Medicine,
685 W. Baltimore St.,
Baltimore, MD 21201, USA

Introduction

Ischemia is the result of cessation of perfusion and the root cause of all stroke cases, mostly originating from blockage of an intra-cranial artery by a blood clot (Galasso and Dyck 2007). While a significant cause of death, stroke causes disability more often than morbidity: a third of all stroke survivors are functionally dependent a year afterwards and stroke ranks as the second most frequent cause of dementia, a prominent cause of depression, and the number one cause of epilepsy in the elderly. Despite the introduction of tissue plasminogen activator and other therapies and interventions which have reduced stroke incidence in some areas (Rothwell et al. 2004), effective treatment for many strokes (or even rapid diagnosis and localization of the insult) still remains to be developed.

Several studies of ischemia have indicated that zinc ion plays a role in subsequent neuronal death (Bossy-Wetzel et al. 2004; Jiang et al. 2001) (reviewed in (Galasso and Dyck 2007; Shuttleworth and Weiss 2011), Thompson (2012)). Even nanomolar zinc levels were observed early on to be

toxic to neurons in culture (Yokoyama et al. 1986), and in some systems elevated zinc induces apoptosis (Telford and Fraker 1995; Aizenman et al. 2000; Bozym et al. 2010), whereas other investigators have found that apoptosis induced by addition of zinc chelators can sometimes be rescued by zinc addition (Waring et al. 1990; Zalewski et al. 1991; Hyun et al. 2001). Furthermore, several investigators have observed substantial increases in extracellular and intracellular free zinc concentrations following ischemia and ischemic-like events (Suh et al. 2000; Wei et al. 2004; Zeng et al. 2005; Frederickson et al. 2006; Stork and Li 2006) and degenerating neurons have been reported to exhibit an increase in intracellular zinc (Tonder et al. 1990; Koh et al. 1996; Aizenman et al. 2000; Yin et al. 2002). Some have proposed that this increase is due to uptake from extracellular sources (notably presynaptic vesicles of a subset of glutamatergic neurons) whereas others believe that the zinc is released from intracellular sources. Another widely observed consequence of ischemia/reperfusion injury is an increase in mitochondrial dysfunction. Ischemic insults result in increased reactive oxygen species (ROS) production, decreased ATP production, activation of mitochondrial permeability transition (MPT), and activation of pro-apoptotic signals (Piantadosi and Zhang 1996; Starkov et al. 2004; Blomgren and Hagberg 2006; Sims and Muyderman 2010; Niizuma et al. 2010). These severe consequences suggest that the integrity of mitochondrial function may well determine overall cellular viability following ischemia/reperfusion. Zinc ions in the sub-micromolar range can directly inhibit key mitochondrial enzymes necessary for energy production such as lipoate-linked dehydrogenases, leading to an increase of ROS production and mitochondrial dysfunction; in addition, zinc ion is a potent inhibitor ($K_i \approx$ nanomolar) of enzymes such as glutathione reductase and thioredoxin reductase, which are key to detoxification of reactive oxygen species (Bogaert et al. 1994; Gazaryan et al. 2002; Gazaryan et al. 2007). Therefore, it would not be surprising if mitochondrial free zinc is under tight control and kept at low concentrations. Consequently, any increase in mitochondrial free zinc, following ischemia/reperfusion, could play a substantial role in the increase in ROS production, decrease in ATP production, and overall mitochondrial dysfunction associated with ischemia/reperfusion injury.

Previously, we described an excitation ratiometric fluorescence biosensor for zinc ion based on human carbonic anhydrase II that exhibits very high affinity and selectivity, and may be inserted into or expressed in cells (Fierke and Thompson 2001; Thompson et al. 2002a; Bozym et al. 2008; Hurst et al. 2010; Wang et al. 2011). The excitation ratiometric feature avoids common artifacts which compromise accuracy when using simple intensity-based fluorescent indicators in cells, due to variations in cell thickness or loading, washout, or photobleaching, while being usable in

ordinary fluorescence microscopes. Moreover, the long emission wavelength of the biosensor (~ 617 nm in this example) avoids the need to correct for autofluorescence of the cells or media. We have demonstrated that the selectivity, sensitivity, and response speed of the biosensor can all be improved by subtle mutagenesis of the protein (Huang et al. 1996; Hunt et al. 1999; McCall and Fierke 2004). Our Förster resonance energy transfer (FRET) based system thus allows for free zinc measurements at picomolar levels (Thompson et al. 2002a) in live resting cells at physiological levels (Bozym et al. 2006). “Free zinc” (or labile zinc) in the cell includes dissolved zinc ion species bound with weak, rapidly exchangeable ligands such as water or chloride ion (Stumm and Morgan 1996); in cells, most zinc ion is tightly bound by proteins, peptides and amino acids (Prasad and Oberleas 1968; Peck and Ray 1971; Krezel and Maret 2006), and is essentially undetected by most indicators, including our biosensor.

The zinc biosensor can be readily inserted into cells via a TAT protein transduction domain (Bozym et al. 2006), but to study mitochondrial zinc we devised an expressible CA II-fluorescent protein fusion which may be targeted to the organelle using the mitochondrial localization sequence of subunit 8 of cytochrome oxidase (Complex IV) in the electron transport chain. We demonstrate that the probe localizes in the mitochondrial matrix, where we can image the free zinc concentration with picomolar affinity. While others have described fluorescent indicators that respond to zinc and localize in the mitochondrion (Sensi et al. 2003; Masanta et al. 2011), as well as expressible (Dittmer et al. 2009) and ratiometric indicators (Caporale et al. 2009), our biosensor has sufficient sensitivity and selectivity to measure mitochondrial matrix zinc at resting levels. The biosensor was calibrated in situ in isolated mitochondria and its matrix location confirmed. Several lines of evidence described above suggest that under ischemic conditions free zinc levels are elevated and contribute to cellular damage; we have now quantitatively imaged free zinc levels in an in vitro ischemia/reperfusion model following the insult. PC-12 cells were subjected to an oxygen glucose deprivation (OGD) model of ischemia (Danilov and Fiskum 2008) which resulted in approximately 50 % cell death 24 h after the end of OGD and their cytoplasmic and mitochondrial free zinc levels were imaged as a function of time following the insult.

Methods

Reagents and materials

The pDsRed2-Mito vector was obtained from Clontech (Mountain View, CA). Lipofectamine 2000, OPTI-MEM media, Neurobasal-A media, B-27 supplement and MitoTracker

Green FM were purchased from Invitrogen (Carlsbad, CA). Digitonin was from Sigma-Aldrich (St. Louis, MO). Goat *anti*-Mouse IgG Dylight_{TM} 405 Conjugated was from Pierce Biotechnology (Rockford, IL). Mouse *anti*-TOM20 antibody was from BD Bioscience (San Jose, CA). Rabbit *anti*-COX IV monoclonal antibody (Alexa Fluor 488 Conjugated) was from Cell Signaling Technology (Danvers, MA). FCCP (carbonyl cyanide 4-(trifluoromethoxy)phenylhydrazone) was from Enzo Life Sciences (Plymouth Meeting, PA). PC12 cells were obtained from ATCC (Manassas, VA). FluoroDish 35 mm glass bottomed tissue culture dishes were from WPI Inc. (Sarasota, FL). All other chemicals and reagents were obtained from common commercial sources.

Construction, expression, and labeling of TAT-H36C-CA cytoplasmic sensor

This sensor was chosen to measure cytoplasmic zinc concentrations because it exhibits almost 20-fold weaker zinc affinity ($K_D=70$ pM) than the wild type (Bozym et al. 2006), extending upward the dynamic range of zinc concentration which it can measure accurately. The construction and fluorescent labeling of the TAT conjugated H36C variant of carbonic anhydrase were as previously described (Bozym et al. 2006). There was no further modification of the probe in this study.

Construction of the CA-DsRed mitochondrial sensor

The gene for human carbonic anhydrase II (CA) was amplified using PCR with the following primers: sense primer, 5'-CCGCGCGCCAAAGATCCATTCGTTGATGGCCCATCACTGGGGGTACGGC-3', containing a BssHII site, and antisense primer, 5'-CATCGGAATCCCCACCTTTGAAGGAAGCTTTG-3'. This CA gene was cloned into the BssHII and blunted BamHI sites of the pDsRed2-Mito vector. The pDsRed2-Mito/CA vector contains the CA gene in frame with an upstream mitochondrial localization signal sequence (from subunit VIII of human cytochrome c oxidase) and a downstream DsRed2 gene. The insert and the expression vector were sequenced to ensure no deleterious mutations had occurred.

Cell culture

PC12 cells were cultured in Neurobasal-A media minus Phenol Red, supplemented with 2 % B-27 supplement, 0.5 mL Glutamine, and 1 % Penicillin-Streptomycin. Prior to experiments PC12 cells were re-plated on 35 mm glass bottomed culture dishes (World Precision) to allow for better imaging.

Cellular transfection

PC12 cells were transfected according to the Lipofectamine 2000 manufacturer's protocol (Invitrogen), using 10 μ g of DNA per 35 mm culture dish. Cells were transfected for 24 h and were used for imaging following another 24 h period.

Mitochondria isolation

Approximately 7.2×10^8 transfected cells were removed from T-75 flasks by trypsinization and quickly resuspended in 10 ml of Dulbecco's phosphate buffered saline (DPBS) containing 5 mg of trypsin inhibitor (0.5 mg/ml). The cells were further diluted by again adding 10 ml of DPBS and centrifuged at 800g for 5 min. The supernatant was removed and cells were resuspended in ice-cold MS buffer (225 mM mannitol, 75 mM sucrose, 5 mM HEPES, 1 mg/ml fatty acid free BSA, pH 7.4) containing 1 mM EGTA and centrifuged at 1,000g for 5 min. The supernatant was again aspirated and cells were resuspended in ice-cold MS buffer+EGTA. The cellular membrane was then disrupted by the addition of 14.4 μ l digitonin from a 10 % stock solution to the cells, while stirring for 30 s. Additional MS buffer and EGTA were then added to the cells to dilute out the digitonin and the cells were centrifuged at 4,000g for 3 min at 4 °C. The supernatant was aspirated and the loose pellet was poured into a Dounce homogenizer. The pellet was homogenized in MS buffer+EGTA, then centrifuged at 2000g for 5 min at 4 °C. After centrifugation the top layer of the supernatant was collected and centrifuged at 14,000g for 10 min at 4 °C. The resulting brown mitochondrial pellet was collected and resuspended in MS buffer+EGTA and centrifuged again at 14,000g for 10 min at 4 °C. The supernatant was then aspirated and the mitochondrial pellet was resuspended in a single drop of MS buffer without EGTA.

Sensor calibration in isolated mitochondria

Approximately 0.45 ng of isolated mitochondria were plated in a 35 mm glass bottomed dish and incubated at 37 °C, 5 % CO₂ for 20 min. The mitochondria were then incubated with 250 μ l each of a series of artificial cerebrospinal fluid (ACSF) zinc buffers pH 7.4, having concentrations of free zinc calculated by MINEQL (Environmental Research Software, Hallowell, ME), for 40 min at 37 °C, 5 % CO₂. After 40 min of incubation 1 μ M of dapoxyl sulfonamide was added to the mitochondria, which were then allowed to incubate for an additional 20 min at 37 °C, 5 % CO₂, prior to imaging.

Sensor co-localization with mitochondria

To establish co-location of the sensor with the mitochondria, cells expressing the sensor were also stained with a

mitochondrion-specific fluorophore, Mitotracker Green. PC12 cells were transfected as described above, and a fraction was plated on 35 mm glass bottomed dishes while the other fraction underwent mitochondrial isolation also as described above. Transfected PC12 cells were incubated with $0.02 \mu\text{M}$ Mitotracker[®] Green FM (Invitrogen) for 30 min at 37°C , 5 % CO_2 , and then washed and imaged using a Nikon Eclipse TE-300 epifluorescence microscope (see below) at excitation and emission wavelengths of 540nm and 617nm , respectively (DsRed2 fluorescence), and also with 490nm excitation and 516nm emission (Mitotracker[®] fluorescence). Images were then merged in IPLab software (Scanalytics, Inc.) to determine co-localization.

Sensor localization within mitochondria

The sensor position with respect to the mitochondrial outer membrane and inner membrane was established by fluorescence colocalization using fluorescent antibodies that recognize proteins specific to the compartments. Isolated mitochondria were split into two groups: one group underwent inner mitochondrial membrane isolation while the other was left untreated. The inner mitochondrial membrane was isolated by dilution with equal parts of a 2 % digitonin stock in 0.25M sucrose solution, which effectively solubilizes the cholesterol-containing outer membrane without solubilizing the cholesterol-poor inner membrane. The resulting inner membrane vesicles (mitoplasts) were stirred gently on ice for 15 min, and then further diluted with 3 volumes of 0.25M sucrose. The mitochondria were then centrifuged for 10 min at $9500g$. The supernatant was removed and saved, and the pellet containing the matrix confined by the inner membrane was resuspended in the sucrose solution and centrifuged for 10 min at $9500g$. Again, the supernatant was removed and combined with the prior fraction, and the pellet was resuspended in a small amount of sucrose solution. Intact mitochondria and inner membrane fractions were each plated on a 35mm glass bottomed dish, and then incubated at 37°C , 5 % CO_2 for 30 min. $2 \mu\text{l}$ of Goat Anti Mouse IgG conjugated with Dylight_{TM} 405 and $2 \mu\text{l}$ of Mouse Anti-TOM20 antibody specific for the outer membrane protein TOM20 were diluted in 1ml of MS Buffer and 1ml of 0.25M sucrose. $10 \mu\text{l}$ of Alexa Fluor 488-conjugated rabbit mAb specific for the COX IV protein of the mitochondrial inner membrane was diluted in 1ml of MS Buffer and 1ml of 0.25M sucrose. An aliquot of $200 \mu\text{l}$ of each antibody solution was added to both the isolated mitochondria and inner mitochondrial membrane fractions, which were then incubated for 30 min at 37°C , 5 % CO_2 . Images were taken at corresponding wavelengths for DsRed2, Dylight 405, and Alexa Fluor 488 and merged as above to determine co-localization.

To determine if the sensor was localized within the mitochondrial matrix, the pH sensitivity of the sensor was utilized: addition of the proton ionophore/uncoupler carbonylcyanide-*p*-trifluoromethoxyphenylhydrazone (FCCP) to respiring mitochondria results in a rapid decline in mitochondrial matrix pH which correspondingly reduces the zinc-binding affinity of the sensor (Lindskog et al. 1971). Thus, if the sensor exhibits an abrupt drop in ratio following FCCP treatment, it most likely resides in the matrix since the pH of the intermembrane space or the extramitochondrial milieu is unaffected by respiratory uncoupling. Transfected PC12 cells were also incubated for 5 min with $1 \mu\text{M}$ Dapoxyl sulfonamide at 37°C , 5 % CO_2 before $0.5 \mu\text{M}$ FCCP dissolved in DMSO was added. Cells were incubated for an additional 15 min and then imaged at Excitation 365nm /Emission 617nm and Excitation 540nm /Emission 617nm . A ratio of Ex 365nm /Ex 540nm was calculated using IPLab software (Scanalytics Inc.) to determine the effect of FCCP on zinc binding.

Hypoxia and hypoglycemia study

As a model of ischemia and reperfusion, cultured PC-12 cells were subjected to oxygen and glucose deprivation, followed by incubation in 95 %/5 % air/ CO_2 (Danilov and Fiskum 2008). For mitochondrial OGD imaging, PC12 cells were transfected as described above. Two days prior to the experiments a zinc- and glucose-free artificial cerebrospinal fluid (ACSF), pH 7.4 (0.142M NaCl, 0.005M KCl, 0.001M $\text{MgCl}_2 \cdot 6\text{H}_2\text{O}$, 0.002M $\text{CaCl}_2 \cdot 4\text{H}_2\text{O}$, 0.01M HEPES, and $50 \mu\text{M}$ Bicine) was deoxygenated in an Anaerobic System (Forma Scientific, Model #1025) in 85 % N_2 , 10 % H_2 , 5 % CO_2 with approximately one ppb oxygen. On the day of the experiment PC-12 cells plated on 35mm glass bottomed culture dishes were placed in the anaerobic chamber and Neurobasal-A medium was replaced with the deoxygenated zinc- and glucose-free ACSF (Jiang et al. 2006). The cells were then incubated in the chamber for 3 h. Following the incubation, the cells were washed three times, supplemented Neurobasal-A media (containing glucose) was added back to the cells, and they were then incubated for 24 h at 37°C , 95 % air, 5 % CO_2 .

Cell staining

For intracellular imaging, cells were stained by immersion in a zinc-free isotonic medium (Thompson et al. 2000) containing $1 \mu\text{M}$ Alexa Fluor 594-labeled apo-TAT-H36C-CA and $1 \mu\text{M}$ Dapoxyl sulfonamide (dissolved in DMSO) and imaged as described previously by Bozym et al. (2006). For mitochondrial imaging of transfected cells only $1 \mu\text{M}$ dapoxyl sulfonamide was added to the cells, which were then incubated for 25 min at 37°C in 5 % CO_2 . Cells were

imaged without washing, in the presence of $1\ \mu\text{M}$ Dapoxyl sulfonamide. Cytosolic zinc images were performed in the presence of Trypan blue staining to ensure that only viable cells were imaged.

Cell imaging

Cells were photographed using a Nikon Eclipse TE300 epifluorescence microscope with a D540/25X or a D360/40X excitation filter, 570DCXRU dichroic and D630-30 M barrier filter (all Omega), through a Nikon Plan Fluor 100x oil/1.3 NA objective with a Cooke Sensicam QE cooled CCD camera. Images were captured using IPLab software (Scanalytics, Inc.). To determine the fluorescence ratio for cells, the images for each excitation ratio were divided and a ratio image was created. Pixels corresponding to the sensor emission (with green excitation) were selected by IPLab software based on relative intensity. The same pixels were then selected in the ratio image to calculate the mean ratio for each image.

Results and discussion

Mitochondria probe calibration and localization

Calibration of the sensor under conditions as similar to the living cell as possible is important to obtain accurate results. While in situ calibration is frequently done in cells for calcium indicators, it has been demonstrated in several cell lines that free zinc ion concentrations above or below a narrow concentration band are toxic (Bozym et al. 2010), and thus such a calibration would be difficult. In view of the widespread use of isolated mitochondria for studying their function, calibrating the sensors in isolated mitochondria was thus most appropriate for this purpose. A plot of the calibration curve of the sensor in unenergized isolated mitochondria in zinc-buffered saline reveals an apparent K_d of $0.15 \pm 0.05\ \text{pM}$ (Fig. 1). This is lower than the K_d of wtCA-DsRed2 in pH 7.4 zinc-saline buffers ($17\ \text{pM}$) and substantially lower than the K_d reported previously (Bozym et al. 2006) for the TAT tagged H36C-AF594-CA probe used for imaging intracellular zinc. The lower K_d is attributable to the high solute levels in the mitochondrion and correspondingly low water activity, which would shift the zinc ion dissociation equilibrium towards the bound rather than the (hydrated) free ion. The exact values of the ratios can vary with the quantum yield of the bound Dapoxyl sulfonamide, which in turn may vary with the altered conditions inside the mitochondrion. The dynamic range of the mitochondrial sensor, wherein the zinc level can be measured with reasonable accuracy, extends roughly from $0.02\ \text{pM}$ to $2\ \text{pM}$.




Fig. 1 Calibration curve for apoCA-DsRed2 expressed in isolated transfected mitochondria (closed squares). The best fit curve shows tighter binding for zinc in isolated mitochondria ($K_d=0.2\ \text{pM}$) compared with ordinary buffers at physiological pH and ionic strength ($K_d=17\ \text{pM}$). The dynamic range for the mitochondrial targeted probe is roughly from $0.02\ \text{pM}$ to $2\ \text{pM}$

We also confirmed that the expressible mitochondria sensor was correctly targeted, by incubating transfected PC12 cells with Mitotracker Green FM, a dye that selectively localizes to the mitochondria. Merged images of both DsRed2 (red fluorescence) and Mitotracker emission (green fluorescence) reveal their co-localization in the mitochondria by yellow color (Fig. 2a). While the co-localization with Mitotracker indicates that the CA-DsRed2 sensor is expressed on or in the mitochondria of transfected cells, standard fluorescence microscopy has insufficient resolution to determine whether the sensor is located within the matrix, the intermembrane space, or bound to the inner or outer membranes. To determine where it is expressed, isolated intact mitochondria and isolated mitochondria with the outer membrane removed by digitonin treatment were probed with fluorescent-labeled antibodies specific for proteins present on the inner (COX IV, a cytochrome oxidase subunit) and outer (Tom20, translocase of outer mitochondrial membrane 20) mitochondrial membranes. Incubations with the TOM20 and COX IV antibodies reveal that the CA-DsRed2 probe is being expressed in the inner membrane or matrix of the mitochondria in transfected cells (Fig. 2b), since the DsRed2 fluorescence of the sensor colocalized with the fluorescence of the anti-TOM20 antibody in the absence (but not the presence) of digitonin treatment, and with the fluorescence of the anti-COX IV antibody following digitonin treatment and centrifugation. To test if the sensor is expressed on the matrix side of the inner mitochondrial membrane as expected, rather than the outer surface of




Fig. 2 Co-localization of Mito-CA-DsRed2. **a** Co-localization of the mitochondrial targeted probe (pseudocolored red) and the MitoTracker probe (pseudocolored green). The yellow color in the merged image indicates that probe successfully targets the mitochondria of transfected cells. **b** Co-localization of the mitochondrial target probe and antibodies specific for the outer mitochondrial membrane (TOM20) and

inner mitochondrial membrane (COX IV). Co-localization experiments were performed on both whole isolated mitochondria and inner mitochondrial membrane/matrix fractions. Incubation with the antibodies suggests that the Mito-CA-DsRed2 sensor is expressed in the inner mitochondrial membrane/matrix of transfected cells


the inner membrane, we observed the effect of FCCP on the reported ratio. By acting as a proton ionophore which uncouples oxidative phosphorylation from ATP synthesis by creating a “proton leak” back into the matrix, FCCP causes a rapid drop in intra-mitochondrial pH shortly after administration. The drop in pH will cause a decrease in the sensor zinc affinity (Lindskog et al. 1971), which will be observed as a decrease in the fluorescence ratio. The observed significant ($p \leq 0.05$) decrease in ratio of excitation 365 nm/excitation 540 nm in cells incubated with 0.5 μM FCCP for 15 min (0.48 ± 0.22), as compared with control cells (0.63 ± 0.11) indicates the sensor is imaging mitochondrial matrix zinc. This is not surprising since the targeting sequence used comes from subunit VIII of cytochrome c oxidase, a matrix protein.

Intracellular and intra-mitochondrial zinc imaging

Previously we had reported imaging intracellular zinc in PC12 cells under physiological conditions using a ratiometric

indicator. From the images it was determined that the average intracellular concentration of labile zinc was around 10 pM, which is somewhat less than reported by other laboratories (Krezel and Maret 2006; Haase et al. 2006; Haase et al. 2008; Rudolf and Červinka 2010) (summarized in Haase and Rink’s excellent review 2011) Specifically, these investigators measured levels between 0.17 and 1 nanomolar free zinc in various cell types (including PC-12 cells) using the fluorescent indicator FluoZin-3. We note that FluoZin-3 is not ratiometric and exhibits a K_D of 15 nM (Sensi et al. 2003), which makes accurate determination of subnanomolar levels in cells by microscopy challenging. Other workers obtained results in the subnanomolar range with Fura-2 (Atar et al. 1995; Ayaz and Turan 2006) and nanomolar range with Mag-Fura-5 (Sensi et al. 1997), but potential competition from endogenous Ca and Mg, respectively, complicates interpretation when using these indicators (Thompson et al. 2002b). Merckx’s group points out (Dongen et al. 2007) that 10 pM is near the bottom of the dynamic range with that CA-based sensor and


Fig. 3 Pseudocolored Ratio Images of Mitochondrial Free [Zn] Following OGD. Physiological conditions (*top left*), 1 h after OGD (*top right*), 2 h after OGD (*bottom left*), 14 h after OGD (*bottom right*). The images clearly show an increase in free zinc in the mitochondria at 1 h after OGD, and a return to physiological levels shortly thereafter



thus the measurement cannot be very precise; while this critique is correct, it is equally clear from this earlier measurement that the free zinc concentration in the cytoplasm of these cells is not 170 pM since this concentration should nearly saturate the wild-type CA-based cytoplasmic sensor and this is not observed using either the TAT-tagged Alexa Fluor 594-

labeled wild type apo CA or E117A CA in eukaryotic PC-12 cells, Chinese hamster ovary cells (Bozym et al. 2006), nor retinal pigmented epithelial ARPE-19 cells (McCranor and Thompson, in preparation), nor the expressible DsRed2-apoCA (wild type) (Bozym and Thompson, unpublished results). Most recently, Qin et al. (2011) measured free zinc

Fig. 4 Changes in cytoplasmic free zinc following OGD. Pseudocolored ratio images following 3 h OGD insult in PC12 cells. Physiological Conditions (*top left*), 1 h after OGD (*top right*), 14 h after OGD (*bottom left*), 24 h after OGD (*bottom right*). Images indicate that intracellular zinc decreases after OGD before increasing gradually over a 24 h time period



concentrations in the Golgi and endoplasmic reticulum compartments of HeLa cells using a targeted expressible sensor and found significantly lower levels (0.6 and 0.9 pM, respectively) in these organelles. Wang et al. (2011) find somewhat higher (20–30 pM) resting levels in *E. coli* using a similar CA-based sensor, and the level varies with extracellular zinc concentration. It does not seem unreasonable that different cell types under varying conditions may in fact have different free zinc concentrations. In this study we imaged intra-mitochondrial zinc using our expressible Mito-CA-DsRed2 sensor to determine the concentration of mitochondrial labile zinc. The sensor reports that, in the mitochondria of PC12 cells under physiological conditions, the average concentration of labile zinc is around 0.15 pM (Fig. 3 upper left).

This number does not necessarily indicate that there is less total zinc in the mitochondrion, rather, it more likely indicates that the matrix contains substantial concentrations of zinc ligands which strongly buffer the free zinc. Note that there is quite substantial variability apparent within even individual cells; it is unclear whether this represents variability in the free zinc concentration, the population of zinc ligands, or both.

Zinc imaging following OGD

In order to mimic the effects of ischemia/reperfusion injury in vitro, an OGD model was employed. PC12 cells were subjected to 3 h of OGD to model ischemia, followed by an additional incubation period (1 h, 2 h, 14 h, or 24 h) under normal conditions (37 °C, 95 % air, 5 % CO₂) to model reperfusion. An incubation of 24 h after 3 h OGD resulted in approximately 50 % dead cells as judged by Live/Dead cell staining (Danilov and Fiskum 2008) (data not shown). Zinc images with the TAT-H36C-CA probe reveal a significant increase in cytoplasmic free zinc 24 h after OGD (>100 pM) as compared to control cells (10 pM) (Fig. 4 and 5). Interestingly, this increase is gradual and follows a period of significantly decreased cytosolic zinc at 1 h after OGD (0.9 pM) (Fig. 4). Again, the cells show increased variability in cytoplasmic zinc levels following OGD as well as (in some cases) reduced nuclear free zinc (Fig. 4, bottom right). At the same time cytosolic zinc is depressed, mitochondrial matrix zinc levels show a substantial increase (to >10 pM) over control levels (0.2 pM) in the first hour after OGD cessation (Fig. 3 and 5); the actual level is not measured very precisely since it is near the top of the sensor dynamic range. However, the surge in mitochondrial free zinc is short lived and by 2 h after OGD it was back down around physiological levels. At later time points, 14 h and 24 h after OGD, a slight increase in mitochondrial zinc is observed, but these increases are not significant and may be due to increased prevalence of non-viable cells.




Fig. 5 Cytosolic and mitochondrial matrix free zinc concentrations following OGD. The average ratio of Ex365nm/Ex540nm was calculated for each cell (left scale), and from this ratio the zinc concentration could be determined using the calibration curve (right scale). After OGD there is a decrease in cytosolic zinc followed by a gradual increase. Mitochondrial matrix zinc increases significantly 1 h after OGD and then returns to near physiological levels. The increase in matrix zinc occurs at the same time as the decrease in cytosolic zinc is observed. Error bars represent the standard error of the mean. $n_{\text{cytosol}}=70-90$ cells, $n_{\text{mito}}=35-45$ cells, * $p < 0.05$, # $p < 0.01$

Discussion

We have developed an expressible ratiometric carbonic anhydrase-based fluorescent indicator system that allows us to quantitatively image free zinc ion in the cytoplasm and specifically in the mitochondrion. We found that the cytoplasmic biosensor gave similar results to those we had previously obtained in other cell types (Bozym et al. 2006). Using this approach we found that the average mitochondrial labile zinc in PC12 cells is in the range of 0.15 pM. This is less than the levels observed in the cytosol, but perhaps unsurprising considering the deleterious effects free zinc can have on mitochondrial function (Jiang et al. 2001; Gazaryan et al. 2002; Gazaryan et al. 2007). We used the biosensors to quantitatively image the changes in intracellular and intra-mitochondrial zinc concentrations in undifferentiated PC12 cells subjected to oxygen + glucose deprivation as a model of ischemia/reperfusion. In both cases the increase in labile zinc was substantial, although at different time points in the study. Over the long term cytosolic zinc is increased in viable cells following OGD, but during the short term (1 h after OGD) cytosolic zinc levels are depressed. At one hour mitochondrial matrix zinc levels are increased substantially (>10 pM), although this measurement is near the top of our

dynamic range. The increase is brief, however, and matrix zinc levels are back nearly at physiological levels by 2 h after OGD.

What is the origin of the zinc changes observed in the cytoplasm and mitochondria? In neuronal models of excitotoxicity, zinc ion can enter cells from the extracellular space through NMDA receptors or AMPA/kainite channels (Sensi et al. 1999; Weiss and Sensi 2000); in this case, while there is higher external free zinc in the medium (≈ 5 nanomolar), there is no glutamate or glutamate analog agonist present to open such channels, so it seems unlikely the cytoplasmic changes occur as a result of zinc movement across the plasma membrane. Zinc can be released from intracellular ligands by at least two mechanisms which are likely to occur during ischemia and reperfusion. The first is oxidation of sulfur ligands on molecules like glutathione, cysteine, and proteins like metallothionein and zinc fingers that bind zinc using cysteine side chains, particularly oxidation of thiols to form disulfides. Oxidation of glutathione during ischemia has been observed (reviewed in Soane et al. 2010) and helps quench reactive oxygen species that are present. The second is release of zinc from histidinyl imidazole ligands by protonation due to acidosis of the tissue. The acidosis arises from lactate production from pyruvate under anaerobic conditions; the brain pH declines as low as 6.0 in models of global ischemia (Fiskum et al. 1999), which protonates histidine residues, reducing their affinity for zinc ions. However, in our model of adherent PC-12 cells (essentially a monolayer) in excess buffered medium, there is likely little change in the pH, so we do not believe this mechanism is important in this case. Finally, we note that several groups have observed release of granular zinc in hippocampal slice preparations following ischemia; since the chemical makeup of the granules remains to be elucidated, the mechanism for the release is unclear. Although the PC-12 cell line is widely used to model neurons (and was chosen in part for this reason), they contain no granular zinc. Thus we conclude that the changes in free zinc ion in our system are largely due to oxidation/reduction of thiol zinc ligands. While we would propose that the elevated levels of zinc in the mitochondrion immediately after OGD cessation are due to release from GSH converted to GSSG and the subsequent decline due to re-reduction of the glutathione, the reasons for the subsequent (modest) increase above normal are unclear. By comparison, there is no ready explanation for the drop in cytoplasmic free zinc following cessation of the insult, nor the steady, significant increase thereafter. It may be that slower process(es) initiated during OGD are operating and result in a release of zinc from intracellular sites. One possibility is that apoptosis (a process that takes hours) initiated during ischemia has resulted in proteolysis of intracellular zinc-binding proteins, releasing bound zinc; we and others have found apoptosis initiated by agents such as

staurosporine results in elevated free zinc in cells prior to their death (results not shown).

These data support other studies referred to above which have observed increases in zinc following ischemia reperfusion injury. Whether the observed increase in mitochondrial matrix zinc is due to an uptake of zinc from the cytosol or a release of zinc from mitochondrial ligands is not known at this time. The results indicate that dramatic changes in zinc homeostasis occur as soon as 1 h after oxygen/glucose deprivation, with differing effects in the cytosol and mitochondria. These changes in zinc concentrations can have a serious effect on overall cell viability, as others have observed, and further implicate zinc as potential cause of cell death following ischemia/reperfusion injury.

Acknowledgments The authors gratefully acknowledge the National Institutes of Health (NIBIB RO1 EB03924 (R.B.T., B.J.M., R.A.B., C.A.F.); PO1 HD16596-25 (G.F., L.H.); NINDS RO1 NS064978 (B.M.P.) and the Susan G. Komen Breast Cancer Foundation (PDF 0601045 (M.I.V.) for their support. The fluorescence sensing technology was originally developed with support from the Office of Naval Research.

References

- Aizenman E, Stout AK, Hartnett KA, Dinely KE, McLaughlin B, Reynolds IJ (2000) Induction of neuronal apoptosis by thiol oxidation: putative role of intracellular zinc release. *J Neurochem* 75(5):1878–1889
- Atar D, Backx PH, Appel MM, Gao WD, Marban E (1995) Excitation-transcription coupling mediated by zinc influx through voltage-dependent calcium channels. *J Biol Chem* 270(6):2473–2477. doi:10.1074/jbc.270.6.2473
- Ayaz M, Turan B (2006) Selenium prevents diabetes-induced alterations in $[Zn^{2+}]_i$ and metallothionein level of rat heart via restoration of cell redox cycle. *Am J Physiol Heart Circ Physiol* 290(3):H1071–H1080. doi:10.1152/ajpheart.00754.2005
- Blomgren H, Hagberg H (2006) Free radicals, mitochondria, and hypoxia-ischemia in the developing brain. *Free Radic Biol Med* 40:388–397
- Bogaert YE, Rosenthal RE, Fiskum G (1994) Postischemic inhibition of cerebral cortex pyruvate dehydrogenase. *Free Radic Biol Med* 16(6):811–820
- Bossy-Wetzel E, Talantova MV, Lee WD, Scholzke MN, Harrop A, Mathews E et al (2004) Crosstalk between nitric oxide and zinc pathways to neuronal cell death involving mitochondrial dysfunction and p38-activated K^+ channels. *Neuron* 41:351–365
- Bozym RA, Thompson RB, Stoddard AK, Fierke CA (2006) Measuring picomolar intracellular exchangeable zinc in PC-12 cells using a ratiometric fluorescence biosensor. *ACS Chem Biol* 1(2):103–111
- Bozym R, Hurst TK, Westerberg N, Stoddard A, Fierke CA, Frederickson CJ et al (2008) Determination of zinc using carbonic anhydrase-based fluorescence biosensors. In: Brand L, Johnson M (eds) *Fluorescence Spectroscopy* (Vol. 450, *Methods in Enzymology*). Academic, San Diego, pp 287–309
- Bozym RA, Chimienti F, Giblin LJ, Gross GW, Korichneva I, Li Y et al (2010) Free zinc outside a narrow concentration range is toxic to a variety of cells in vitro. *Experimental Biology and Medicine* 235:741–750. doi:10.1258/ebm.2010.009258

- Caporale T, Ciavardelli D, Di Ilio C, Lanuti P, Drago D, Sensi SL (2009) Ratiometric-pericam-mt, a novel tool to evaluate intramitochondrial zinc. *Exp Neurol* 218(2):228
- Danilov CA, Fiskum G (2008) Hyperoxia promotes astrocyte cell death after oxygen and glucose deprivation. *Glia* 56(7):801–808. doi:10.1002/glia.20655
- Dittmer PJ, Miranda JG, Gorski JA, Palmer AE (2009) Genetically encoded sensors to elucidate spatial distribution of cellular zinc. *J Biol Chem* 284:16289–16297. doi:10.1074/jbc.M900501200
- Dongen EMWV, Evers TH, Dekkers LM, Meijer EW, Klomp LWJ, Merx M (2007) Variation of linker length in ratiometric fluorescent sensor proteins allows rational tuning of Zn(II) affinity in the picomolar to femtomolar range. *Journal of the American Chemical Society* 129:3494–3495. doi:10.1021/ja069105
- Fierke CA, Thompson RB (2001) Fluorescence-based biosensing of zinc using carbonic anhydrase. *BioMetals* 14:205–222
- Fiskum G, Murphy AN, Beal MF (1999) Mitochondria in neurodegeneration: acute ischemia and chronic neurodegenerative diseases. *J Cereb Blood Flow Metab* 19:351–369
- Frederickson CJ, Giblin LJ, Krezel A, McAdoo DJ, Muelle RN, Zeng Y et al (2006) Concentrations of extracellular free zinc (pZn) in the central nervous system during simple anesthetization, ischemia, and reperfusion. *Exp Neurol* 198:285–293. doi:10.1016/j.expneurol.2005.08.030
- Galasso SL, Dyck RH (2007) The role of zinc in cerebral ischemia. *Molecular Medicine* 13(7–8):380–387. doi:10.2119/2007-00044.Galasso
- Gazaryan I, Krasnikov BF, Ashby GA, Thorneley RNF, Kristal BS, Brown AM (2002) Zinc is a potent inhibitor of thiol oxidoreductase activity and stimulates reactive oxygen species production by lipoamide dehydrogenase. *J Biol Chem* 277(12):10064–10072
- Gazaryan IG, Krasinskaya IP, Kristal BS, Brown AM (2007) Zinc irreversibly damages major enzymes of energy production and antioxidant defense prior to mitochondrial permeability transition. *J Biol Chem* 282:24373–24380. doi:10.1074/jbc.M611376200
- Haase H, Rink L (2011) Zinc signaling. In: Rink L (ed) *Zinc in human health*. IOS, Amsterdam
- Haase H, Hebel S, Engelhardt G, Rink L (2006) Flow cytometric measurement of labile zinc in peripheral blood mononuclear cells. *Anal Biochem* 352(2):222
- Haase H, Ober-Blißbaum JL, Engelhardt G, Hebel S, Heit A, Heine H et al (2008) Zinc signals are essential for lipopolysaccharide-induced signal transduction in monocytes. *J Immunol* 181(9):6491–6502
- Huang C-C, Lesburg CA, Kiefer LL, Fierke CA, Christianson DW (1996) Reversal of the hydrogen bond to zinc ligand histidine-119 dramatically diminishes catalysis and enhances metal equilibration kinetics in carbonic anhydrase II. *Biochemistry* 35(11):3439–3446
- Hunt JA, Ahmed M, Fierke CA (1999) Metal binding specificity in carbonic anhydrase is influenced by conserved hydrophobic amino acids. *Biochemistry* 38:9054–9060
- Hurst TK, Wang D, Thompson RB, Fierke CA (2010) Carbonic anhydrase II-based metal ion sensing: Advances and new perspectives. *Biochimica et Biophysica Acta (BBA) - Proteins & Proteomics* 1804(2):393–403
- Hyun HJ, Sohn JH, Ha DW, Ahn YH, Koh JY, Yoon YH (2001) Depletion of intracellular zinc and copper with TPEN results in apoptosis of cultured human retinal pigment epithelial cells. *Invest Ophthalmol Vis Sci* 42(2):460–465
- Jiang D, Sullivan PG, Sensi SL, Steward O, Weiss JH (2001) Zn²⁺ induces permeability transition pore opening and release of proapoptotic peptides from neuronal mitochondria. *J Biol Chem* 276:47524–47529
- Jiang Z-G, Lu XCM, Nelson V, Yang X, Pan W, Chen R-W et al (2006) A multifunctional cytoprotective agent that reduces neurodegeneration after ischemia. *Proc Natl Acad Sci U S A* 103(5):1581–1586. doi:10.1073/pnas.0510573103
- Koh JY, Suh S-W, Gwag BJ, He YY, Hsu CY, Choi DW (1996) The role of zinc in selective neuronal death after transient global cerebral ischemia. *Science* 272:1013–1016
- Krezel A, Maret W (2006) Zinc-buffering capacity of a eukaryotic cell at physiological pZn. *J Biol Inorg Chem* 11:1049–1062
- Lindskog S, Henderson LE, Kannan KK, Liljas A, Nyman PO, Strandberg B (1971) Carbonic anhydrase. In: Boyer PD (ed) *The enzymes* (Vol. 5, *The Enzymes*). Academic, New York, pp 587–665
- Masanta G, Lim CS, Kim HJ, Han JH, Kim HM, Cho BR (2011) A mitochondrial-targeted two-photon probe for zinc ion. *J Am Chem Soc* 133(15):5698
- McCall KA, Fierke CA (2004) Probing determinants of the metal ion selectivity in carbonic anhydrase using mutagenesis. *Biochemistry* 43:3979–3986
- Niizuma K, Yoshioka H, Chen H, Kim GS, Jung JE, Katsu M et al (2010) Mitochondrial and apoptotic neuronal death signaling pathways in cerebral ischemia. *Biochim Biophys Acta* 1802:92–99
- Peck EJ, Ray WJ (1971) Metal complexes of phosphoglucomutase *in vivo*: alterations induced by insulin. *J Biol Chem* 246(4):1160–1167
- Piantadosi CA, Zhang J (1996) Mitochondrial generation of reactive oxygen species after brain ischemia in the rat. *Stroke* 27:327–332
- Prasad AS, Oberleas D (1968) Zinc in human serum: evidence for an amino acid-bound fraction. *J Lab Clin Med* 1006
- Qin Y, Dittmer PJ, Park JG, Jansen KB, Palmer AE (2011) Measuring steady-state and dynamic endoplasmic reticulum and Golgi Zn²⁺ with genetically encoded sensors. *Proc Natl Acad Sci*. doi:10.1037/pnas.1015686108
- Rothwell PM, Coull AJ, Giles MF, Howard SC, Silver LE, Bull LM et al (2004) Change in stroke incidence, mortality, case-fatality, severity, and risk factors in Oxfordshire, UK from 1981 to 2004 (Oxford Vascular Study). *Lancet* 363(9425):1925
- Rudolf E, Červinka M (2010) Zinc pyrithione induces cellular stress signaling and apoptosis in Hep-2 cervical tumor cells: the role of mitochondria and lysosomes. *BioMetals* 23(2):339–354
- Sensi SL, Canoniero LMT, Yu SP, Ying HS, Koh J-Y, Kerchner GA et al (1997) Measurement of intracellular free zinc in living cortical neurons: routes of entry. *J Neurosci* 17(24):9554–9564
- Sensi SL, Yin HZ, Carriedo SG, Rao SS, Weiss JH (1999) Preferential Zn²⁺ influx through Ca²⁺-permeable AMPA/kainate channels triggers prolonged mitochondrial superoxide production. *Proc Natl Acad Sci* 96:2414–2419
- Sensi SL, Ton-That D, Weiss JH, Rothe A, Gee KR (2003) A new mitochondrial fluorescent zinc sensor. *Cell Calcium* 34(3):281–284
- Shuttleworth CW, Weiss JH (2011) Zinc: new clues to diverse roles in brain ischemia. *Trends Pharmacol Sci* 32(8):480. doi:10.1016/j.tips.2011.04.001/j.tips.2011.04.001
- Sims NR, Muyderman H (2010) Mitochondria, oxidative metabolism and cell death in stroke. *Biochim Biophys Acta* 1802:80–91
- Soane L, Polster BM, Fiskum G (2010) Mitochondrial mechanisms of neural cell death in cerebral ischemia. In: Reed JC, Green D (eds) *Apoptosis: physiology and pathology of cell death*. Cambridge University Press, Cambridge
- Starkov AA, Fiskum G, Chinopoulos C, Lorenzo BJ, Browne SE, Patel MS et al (2004) Mitochondrial alpha-Ketoglutarate Dehydrogenase Complex Generates Reactive Oxygen Species. *J Neurosci* 24(36):7779–7788. doi:10.1523/jneurosci.1899-04.2004
- Stork CJ, Li YV (2006) Intracellular zinc elevation measured with a "calcium-specific" indicator during ischemia and reperfusion in rat hippocampus: a question on calcium overload. *J Neurosci* 26(41):10430–10437. doi:10.1523/JNEUROSCI.1588-06.2006
- Stumm W, Morgan JJ (1996) *Aquatic chemistry: chemical equilibria and rates in natural waters*, 3rd edn. Wiley-Interscience, New York

- Suh SW, Chen JW, Motamedi M, Bell B, Listiak K, Pons NF et al (2000) Evidence that synaptically-released zinc contributes to neuronal injury after traumatic brain injury. *Brain Res* 852:268–273
- Telford W, Fraker PJ (1995) Preferential induction of apoptosis in mouse CD4⁺ CD8⁺ alpha betaTCR10CD3e thymocytes by zinc. *J Cell Physiol* 164:259–270
- Thompson RB (2012) Zinc in stroke: Time for a new approach? In: Li YV, Zhang JH (eds) *Metal Ions in Stroke*. Springer, Berlin, in the press. doi:10.1007/978-1-4419-9663-3_9
- Thompson RB Jr, Whetsell OW, Maliwal BP, Fierke CA, Frederickson CJ (2000) Fluorescence microscopy of stimulated Zn(II) release from organotypic cultures of mammalian hippocampus using a carbonic anhydrase-based biosensor system. *J Neurosci Meth* 96(1):35–45
- Thompson RB, Cramer ML, Bozym R, Fierke CA (2002a) Excitation ratiometric fluorescent biosensor for zinc ion at picomolar levels. *J Biomed Opt* 7(4):555–560
- Thompson RB, Peterson D, Mahoney W, Cramer M, Maliwal BP, Suh SW et al (2002b) Fluorescent zinc indicators for neurobiology. *J Neurosci Meth* 118:63–75
- Tonder N, Johansen FF, Frederickson CJ, Zimmer J, Diemer NH (1990) Possible role of zinc in the selective degeneration of dentate hilar neurons after cerebral ischemia in the adult rat. *Neurosci Lett* 109:247–252
- Wang D, Hurst TK, Thompson RB, Fierke CA (2011) Genetically encoded ratiometric biosensors to measure extracellular exchangeable zinc in *Escherichia coli*. *J Biomed Opt* 16(8):087011. doi:10.1117/1.3613926
- Waring P, Egan M, Braithwaite A, Mullbacher N, Siaarda A (1990) Apoptosis induced in macrophages and T-blasts by the mycotoxin sporidesmin and protection by zinc salts. *Int J Pharmacol* 12:445–457
- Wei G, Hough CJ, Li Y, Sarvey JM (2004) Characterization of extracellular accumulation of Zn²⁺ during ischemia and reperfusion of hippocampus slices in rat. *Neuroscience* 125:867–877
- Weiss JH, Sensi S (2000) Ca²⁺-Zn²⁺ permeable AMPA or kainate receptors: possible key factors in selective neurodegeneration. *Trends Neurosci* 23(8):365–371
- Yin HZ, Sensi SL, Ogoshi F, Weiss JH (2002) Blockade of Ca²⁺-permeable AMPA/kainate channels decreases oxygen-glucose deprivation-induced Zn²⁺ accumulation and neuronal loss in hippocampal pyramidal neurons. *J Neurosci* 22(4):1273–1279
- Yokoyama M, Koh J, Choi DW (1986) Brief exposure to zinc is toxic to cortical neurons. *Neurosci Lett* 71:351–355
- Zalewski P, Forbes IJ, Giannakis C (1991) Physiological role for zinc in prevention of apoptosis (gene-directed death). *Biochem Int* 24:1093–1101
- Zeng H-H, Bozym RA, Rosenthal RE, Fiskum G, Cotto-Cumba C, Westerberg N, et al. (2005) In situ measurement of free zinc in an ischemia model and cell culture using a ratiometric fluorescence-based biosensor. In T. Vo-Dinh, W. S. Grundfest, D. A. Benaron, & G. E. Cohn (Eds.), *SPIE Conference on Advanced Biomedical and Clinical Diagnostic Systems III, San Jose, CA*, (Vol. 5692, pp. 51–59): SPIE