



## Review

# Mitochondrial mechanisms of cell death and neuroprotection in pediatric ischemic and traumatic brain injury

Courtney L. Robertson<sup>a,b</sup>, Susanna Scafidi<sup>a</sup>, Mary C. McKenna<sup>a</sup>, Gary Fiskum<sup>b,c,\*</sup>

<sup>a</sup> Department of Pediatrics, University of Maryland School of Medicine, Baltimore MD, USA

<sup>b</sup> Department of Anesthesiology, University of Maryland School of Medicine, Baltimore MD, USA

<sup>c</sup> Trauma and Anesthesiology Research Center, University of Maryland School of Medicine, Baltimore MD, USA

## ARTICLE INFO

## Article history:

Received 17 January 2009

Revised 23 April 2009

Accepted 28 April 2009

Available online 7 May 2009

## Keywords:

Metabolism

Calcium

Permeability transition pore

poly(ADP-ribose) polymerase

Reactive oxygen species

Pyruvate dehydrogenase

Cytochrome c

Nicotinamide adenine dinucleotide

## ABSTRACT

There are several forms of acute pediatric brain injury, including neonatal asphyxia, pediatric cardiac arrest with global ischemia, and head trauma, that result in devastating, lifelong neurologic impairment. The only clinical intervention that appears neuroprotective is hypothermia initiated soon after the initial injury. Evidence indicates that oxidative stress, mitochondrial dysfunction, and impaired cerebral energy metabolism contribute to the brain cell death that is responsible for much of the poor neurologic outcome from these events. Recent results obtained from both *in vitro* and animal models of neuronal death in the immature brain point toward several molecular mechanisms that are either induced or promoted by oxidative modification of macromolecules, including consumption of cytosolic and mitochondrial NAD<sup>+</sup> by poly-ADP ribose polymerase, opening of the mitochondrial inner membrane permeability transition pore, and inactivation of key, rate-limiting metabolic enzymes, e.g., the pyruvate dehydrogenase complex. In addition, the relative abundance of pro-apoptotic proteins in immature brains and neurons, and particularly within their mitochondria, predisposes these cells to the intrinsic, mitochondrial pathway of apoptosis, mediated by Bax- or Bak-triggered release of proteins into the cytosol through the mitochondrial outer membrane. Based on these pathways of cell dysfunction and death, several approaches toward neuroprotection are being investigated that show promise toward clinical translation. These strategies include minimizing oxidative stress by avoiding unnecessary hyperoxia, promoting aerobic energy metabolism by repletion of NAD<sup>+</sup> and by providing alternative oxidative fuels, e.g., ketone bodies, directly interfering with apoptotic pathways at the mitochondrial level, and pharmacologic induction of antioxidant and anti-inflammatory gene expression.

© 2009 Published by Elsevier Inc.

## Contents

Pediatric hypoxic and traumatic brain injury . . . . .	371
Neuroprotection for pediatric brain injury . . . . .	372
Oxidative stress, mitochondrial dysfunction, and metabolic failure . . . . .	373
Mitochondria and the developing brain . . . . .	374
Mitochondrial pathways for neuroprotection in the immature brain . . . . .	375
Acknowledgments . . . . .	377
References . . . . .	377

## Pediatric hypoxic and traumatic brain injury

Brain injury in the pediatric population can have a great economic and emotional impact on both the child's family and

society. It is important to remember that in addition to regaining lost skills, developing children are expected to rapidly be developing new skills, making the burden of brain injury even greater. Children with severe injury can require an extended amount of chronic care throughout their lifetime. One of the most common causes of permanent neurologic injury in children is perinatal hypoxia-ischemia. Infants sustain asphyxial injury around the time of birth at a rate of 0.2 to 0.4% of all full-term births. The incidence is much

\* Corresponding author. Dept. of Anesthesiology, University of Maryland School of Medicine, 685 W. Baltimore St., Baltimore, MD 21201, USA. Fax: +1 410 706 2550.

E-mail address: [gfishum@anes.umm.edu](mailto:gfishum@anes.umm.edu) (G. Fiskum).

greater in the preterm and low birth weight infant population (MacDonald et al., 1980). Up to 50% of the most severely affected infants will die in the newborn period, and the less severely injured infants can survive with significant, permanent neurologic deficits. These can include cerebral palsy, mental retardation, learning disabilities, epilepsy, deafness and blindness. Importantly, long-term outcome studies have shown that 20% of infants demonstrating moderate encephalopathy secondary to perinatal hypoxia–ischemia have marked neurologic deficits at 3.5 and 8 years of age (Robertson et al., 1989). Furthermore, in the subset of children that did not have marked deficits, up to 1/3 had delays in their academic progress in school.

Another cause of neurologic injury in children is cardiac arrest. There are many etiologies of cardiac arrest in children, but the majority (>60%) involve a period of asphyxia preceding loss of circulation (Manole et al., 2008; Berg et al., 2008). Outcome is relatively poor, as only ~30% of patients have return of spontaneous circulation, and only 12% of these patients survive to hospital discharge (Donoghue et al., 2005). The burden of neurologic impairment is high in survivors, with only ~4% having intact neurologic outcomes (no or mild impairment) (Donoghue et al., 2005).

Trauma is the leading cause of death in children ages 1–18, and traumatic brain injury (TBI) is the major determinant of outcome in these injured children (NIH 1998). Every year in the United States alone, almost 1/2 million children sustain traumatic brain injury, and ~3000 children per year die from these injuries (CDC). Despite recent advances in neurointensive care and reduction in the overall mortality rate (Tilford et al., 2005), the long-term morbidity of severe TBI in childhood remains high. Survivors of pediatric TBI suffer from many long-term physical, cognitive, psychological, and emotional impairments (Yeates et al., 2004; Max et al., 2004; Wechsler et al., 2005; Keenan and Bratton, 2006). For example, one study found that 40% of children with severe TBI had a persistent personality change after injury (Max et al., 2000), and other studies have shown profound declines in long-term academic performance (Ewing-Cobbs et al., 2004a). Importantly, children injured at younger ages (<4–6 years) may be especially vulnerable to declines in intelligence and executive functioning (Ewing-Cobbs et al., 2006), (Ewing-Cobbs et al., 2004b; Anderson et al., 2005). Despite these daunting facts, there are currently no effective neuroprotective therapies available for children with severe head injury. Additional studies elucidating the age-specific mechanisms of secondary injury and recovery are essential, in order to target these mechanisms for effective neuroprotective treatments in pediatric acute brain injury.

### Neuroprotection for pediatric brain injury

Current therapeutic options for pediatric brain injury are limited and essentially focus on supportive care of the injured infant and avoidance of any additional systemic insults such as further hypoxia, hypotension, and secondary ischemia due to cerebral edema. Of note, most of these parameters, with the exception of hypoxia, have age-dependent definitions. For instance, normal blood pressure ranges increase steadily with age, and normal cerebral blood flow (CBF) is higher in children than in infants or teenagers. Peak global CBF is seen at ~3–4 years of age (108 ml/100 g/min), with a slow decline into the teenage years (~70 ml/100 g/min) (Suzuki, 1990). Most normal values for children remain above the normal CBF range for adults (~50 ml/100 g/min), with the exception of infants <6 months of age (~40 ml/100 g/min) (Suzuki, 1990) (Zwieneberg and Muizelaar, 1999).

No pharmacologic or therapeutic interventions, either alone or in combination, have proven effective in improving neurologic outcome of infants and children after asphyxial or traumatic insults. In fact, most medical therapies used in the past to treat hypoxic–

ischemic brain injury, such as steroids, barbiturates and osmotic agents, have proven ineffective (Vannucci, 1990). Clinical trials in pediatric TBI are similarly limited (Natale et al., 2006), and the majority of clinical management guidelines are extrapolated from adult TBI guidelines (Adelson et al., 2003; Jankowitz and Adelson, 2006). Two main therapeutic interventions that have been tested prospectively in head-injured children recently are decompressive craniectomy (Taylor et al., 2001) and hypothermia (Biswas et al., 2002; Hutchison et al., 2008). Early decompressive craniectomy in children reduced intracranial hypertension and showed a trend toward improved long-term outcome, but the study was limited by small numbers ( $n=27$ ) (Taylor et al., 2001). Importantly, the recently completed multi-center randomized control trial on hypothermia for severe pediatric TBI (32.5 °C for 24 h) did not significantly improve neurologic outcome (Hutchison et al., 2008) (see discussion below). Discovery of successful treatment strategies will require continued study of the cellular and subcellular pathophysiological changes that occur in the brain subsequent to hypoxia–ischemia or trauma. The use of animal models of perinatal hypoxia–ischemia or head trauma that shows injury patterns similar to those seen in human newborns will facilitate this process (Vannucci et al. 1999, Prins and Hovda, 2003).

It is now well known that injury to the brain initiated during hypoxia or ischemia continues, and is magnified during the post-resuscitative period. This “reperfusion illness” is multifactorial in etiology and is caused in part by such processes as excitotoxicity, oxidative stress, metabolic failure, inflammation, apoptosis, and others. While many drugs provide neuroprotection in animal models of global cerebral ischemia, prevention of post-ischemic human brain injury through pharmacologic intervention has not yet been achieved (Harukuni and Bhardwaj, 2006). For example, no increased survival or neuroprotection among cardiac arrest (CA) survivors was demonstrated in adults through the use of glucocorticoids (Jastremski et al., 1989) (Jastremski et al., 1989), thiopental (Brain Resuscitation Clinical Trial I Study Group, 1986), or calcium channel antagonists (Brain Resuscitation Clinical Trial II Study Group, 1991). Similarly, several large clinical trials failed to demonstrate a role for high dose epinephrine in human victims of CA. (Goetting and Paradis 1991; Barton and Callahan 1991). Great excitement was generated in 2002, however, when two large clinical trials demonstrated significant improvement in neurologic outcome and reduced mortality in adults treated with moderate hypothermia following resuscitation from CA. (Bernard et al., 2002; Hypothermia Cardiac Arrest Study Group, 2002). This breakthrough helped renew the conviction that lessons learned in the laboratory can be successfully translated into improved long-term resuscitation outcomes following human CA. Pediatric studies of hypothermia to date have focused primarily on neonatal asphyxia and TBI. In neonates, both whole-body cooling (esophageal  $T=33.5$  °C for 72 h) or selective head cooling (rectal  $T=34–35$  °C, head cooling unit with water  $T=8–12$  °C for 72 h) have shown promise in reducing the risk of death or severe disability (Shankaran et al., 2005; Gluckman et al., 2005). Preliminary evidence from clinical trials testing head cooling for neuroprotection after neonatal hypoxic ischemic encephalopathy suggests efficacy (Wyatt et al., 2007; Gunn et al., 2008), and a large multi-center clinical trial testing whole-body cooling on survival and neurologic outcome is in progress (Azzopardi et al., 2008). Initial studies of hypothermia in pediatric TBI (brain  $T=32–33$  °C) have demonstrated that the treatment is relatively safe and improves control of ICP (Biswas et al., 2002; Adelson et al., 2005). In contrast, Hutchison et al. (2008) did not show efficacy of hypothermia (brain  $T=32.5$  °C for 24 h) initiated within 8 h and continued for 24 h after pediatric TBI. In contrast to adult stroke victims, who often do not receive medical attention for many hours after the stroke onset, the vast majority of TBI victims reach a hospital within an hour or two after the

injury and are therefore available for treatment. Together with successfully resuscitated cardiac arrest survivors, TBI victims can receive potentially neuroprotective interventions, e.g., hypothermia, relatively early after the injury, thereby increasing the likelihood of success.

In order for any neuroprotective strategy (including hypothermia) to make an impact, however, the technique must be safe, easy to implement, and widely accepted. Despite successful hypothermia clinical trials, the emergency medicine/critical care community has not yet embraced the use of hypothermia in the clinical arena. Even if widely accepted, one recent editorial suggests that hypothermia should be only one part of a strategy designed to attack different components of the pathophysiologic cascade (Broccard, 2006). We have established that by avoiding unnecessary hyperoxia during the first hour following resuscitation after global cerebral ischemia (cardiac arrest) in adult animals, we significantly reduce oxidative stress, metabolic dysfunction, delayed neuronal death, and both short-term and long-term neurobehavioral impairment (Liu et al., 1998; Richards et al., 2006; Balan et al., 2006; Vereczki et al., 2006; Richards et al., 2007). Comparisons of the effects of hyperoxic and normoxic resuscitation in neonatal rat models of hypoxic ischemia are equivocal. One study found no overall difference in long-term histologic outcome after reperfusion under 100%, 40%, or 21% inspired O<sub>2</sub>; however, there was a trend toward better outcome specifically for male rats using 40% O<sub>2</sub> (Grafe et al., 2008). Another study demonstrated that 100% O<sub>2</sub> after neonatal HI reduces cerebral blood flow (CBF) and, importantly, increases brain superoxide production (Fabian et al., 2008). Importantly, clinical studies comparing room air (21% O<sub>2</sub>) to hyperoxic (100% O<sub>2</sub>) resuscitation in depressed newborns have generally shown equivalent long-term neurologic outcomes between groups, and a recent meta-analysis found a lower mortality rate in room air resuscitated infants (Rabi et al., 2007). In pediatric cardiac arrest (CA), preclinical studies of asphyxial CA in immature rats found increased markers of tissue oxidative injury in rats resuscitated with hyperoxic conditions (Walson et al., 2008). To our knowledge, the effect of hyperoxic versus normoxic resuscitation on outcome after pediatric TBI has not been studied. In adult TBI, hyperoxic resuscitation has produced mixed results (Diringer, 2008). Some studies have shown that hyperoxia improves cerebral metabolism (Tolias et al., 2004; Tisdall et al., 2008), while others suggest it can produce oxidative injury to cellular constituents (Ahn et al., 2008). The role of oxygen following injury to the developing brain is clearly a controversial but clinically very important topic that requires more research using both animal models and clinical trials. While better management of oxygenation to avoid both hypoxia and unnecessary hyperoxia will likely improve clinical outcome, it is unlikely that this intervention alone will provide optimal long-term result. Additional approaches include those that target mitochondrial pathways of necrotic and apoptotic cell death.


### Oxidative stress, mitochondrial dysfunction, and metabolic failure

A large body of evidence indicates that mitochondrial dysfunction plays a critical role in the pathophysiology of ischemic and traumatic brain injury (Blomgren and Hagberg, 2006; Chang et al., 1992; Fiskum et al., 1999; Kuroda et al., 1996; Rizzuto et al., 1992; Starkov et al., 2004). Consequences of mitochondrial dysfunction are numerous and include oxidative stress, loss of cellular Ca<sup>2+</sup> homeostasis, promotion of apoptosis, and metabolic failure.

There are many possible causes of mitochondrial metabolic impairment and most involve oxidative modifications to proteins, lipids, or DNA. Identification of the sites at which oxidative stress impairs respiration can guide the development of counteractive interventions with neuroprotective potential. Complex I of the electron transport chain (ETC), which catalyzes the oxidation of NADH and the reduction of ubiquinone, is particularly sensitive to inhibition by both

oxidative stress and ischemia/reperfusion and is generally considered to be the rate-limiting component of the ETC (Hillered and Ernster, 1983; Rosenthal and Fiskum, 1990; Rosenthal et al., 1987; Sims, 1991). Another cause of impaired ETC activity is the release of cytochrome c through the outer mitochondrial membrane into the cytosol, an event that is also often followed by caspase-dependent apoptosis (Polster et al. 2001). Oxidative stress promotes cytochrome c release by several mechanisms, including those promoting translocation of Bax and Bak to the mitochondrial outer membrane (Fig. 1) (Castino et al. 2007; Perier et al. 2005). These proteins form megapores within the outer membrane when prompted to oligomerize by the binding of BH3 domain only proteins, e.g., tBid, to these proteins or to anti-apoptotic proteins, e.g., Bcl2 or Bclx<sub>L</sub>, that normally block megapore formation by heterodimerizing with Bax and Bak. Although not previously measured, the loss of cytochrome c through such megapores could be responsible for the secondary brain mitochondrial respiratory impairment observed 24 h after neonatal HI, when caspase 3 activation is robust (Puka-Sundvall et al., 2000d).

In addition to impaired ETC activities, oxidative phosphorylation can also be obstructed by inhibition of other mitochondrial enzymes and membrane transporters (Fig. 1). Thus, oxidative inactivation of mitochondrial matrix enzymes, e.g., pyruvate and  $\alpha$ -ketoglutarate dehydrogenases and aconitase, are implicated in metabolic failure (Bogaert et al., 1994; Richards et al., 2006; Vereczki et al., 2006). Evidence also suggests that mitochondrial oxidative stress impairs the adenine nucleotide translocase, necessary for influx of ADP and efflux of ATP (Vesce et al., 2005). While there is little if any information about the effects of acute brain injury on other mitochondrial membrane transporters, one recent study suggests that even transient elevation of intramitochondrial Ca<sup>2+</sup> can stimulate glycolytic lactate production through impairment of net efflux of mitochondrial  $\alpha$ -ketoglutarate and subsequent inhibition of the malate/aspartate shuttle system (Contreras and Satrustegui, 2009). Stimulation of  $\alpha$ -ketoglutarate dehydrogenase by Ca<sup>2+</sup> can reduce the level of  $\alpha$ -ketoglutarate within both brain and heart mitochondria to a level that limits its efflux through the oxoglutarate transporter. The lack of cytosolic  $\alpha$ -ketoglutarate then limits the aspartate transaminase reaction that is necessary for the shuttling of reducing power from glycolysis-generated NADH to the mitochondria. Consequently, lactate dehydrogenase utilizes this NADH to form lactate from pyruvate, thus shunting pyruvate away from



**Fig. 1.** Oxidative stress and mitochondrial metabolic dysfunction. Mechanisms of mitochondrial and metabolic dysfunction following acute brain injury include direct inhibition of the pyruvate dehydrogenase complex (PDHC) by reactive O<sub>2</sub> species (ROS), loss of mitochondrial NAD(H) through the permeability transition pore (PTP), consumption of NAD<sup>+</sup> by poly-ADP ribose polymerase 1 (PARP), and release of cytochrome c (c) through Bax or Bak megapores present in the outer membrane. Respiratory inhibition stimulates lactate dehydrogenase (LDH), resulting in lactic acidosis, which further inhibits oxidative phosphorylation and promotes oxidative stress.

mitochondrial aerobic metabolism. While this scenario is likely to be very transient under physiological conditions, it may persist for some time after cerebral ischemia, head trauma, and other events associated with excitotoxicity, where mitochondria are exposed to and sequester elevated cytosolic  $\text{Ca}^{2+}$  for prolonged periods.

Much interest is also currently focused on the availability of the metabolic cofactor  $\text{NAD}^+$ , necessary for the numerous dehydrogenases present within the mitochondrial matrix (Ying, 2008). Cellular  $\text{NAD}^+$  can be rapidly catabolized by nuclear and possibly mitochondrial poly-ADP-ribose polymerase in response to activation of this enzyme by oxidative DNA modifications and ERK kinase-mediated enzyme phosphorylation (Du et al., 2003; Eliasson et al., 1997; Kauppinen and Swanson, 2007).  $\text{NAD}^+$  in its oxidized or reduced form ( $\text{NADH}$ ) can be lost from the mitochondrial matrix following opening of the inner membrane permeability transition pore (PTP), which results in transmembrane equilibration of small ions and molecules of up to approximately 1500 Da (Fig. 1) (Crompton et al., 2002; Halestrap et al., 2002). The PTP is activated by abnormally high concentrations of  $\text{Ca}^{2+}$  and by oxidative stress (Bernardi et al., 1999). Little is known regarding the effects of brain development on mitochondrial permeability transition. One study provided evidence that the PTP is relatively sensitive to  $\text{Ca}^{2+}$ -induced opening and cyclosporin A-mediated inhibition in mitochondria from immature rats (Eliseev et al., 2007). We found that the relative sensitivity of PTP opening in immature rat brain mitochondria is dependent on the environment, e.g., the presence of ATP and the pH. In fact, the PTP is comparatively very active in immature brain mitochondria at low pH and ATP, conditions that are present during and immediately following hypoxic ischemia (Robertson et al., 2004).

Contribution of PTP opening to ischemic and traumatic brain injury is supported by the neuroprotection observed with PTP inhibitors, e.g., cyclosporins (Alessandri et al., 2002; Hansson et al., 2004; Sullivan et

al., 2002; Uchino et al., 2002), that bind to cyclophilin D, the one well-established protein associated with pore opening. Cyclophilin D knock-out mice are resistant to ischemic brain injury (Schinzel et al., 2005), and to spinal cord axonal damage associated with experimental autoimmune encephalomyelitis, an animal model characterized by extensive inflammation (Forte et al., 2007). These findings taken together with observations that post-ischemic inflammation is accompanied by considerable ROS formation via iNOS and NADPH oxidase (Wang et al., 2007), strongly suggest that the PTP participates in inflammatory neurodegeneration after global cerebral ischemia. Support for involvement of the PTP in neonatal hypoxic ischemia comes from the observations of mitochondrial  $\text{Ca}^{2+}$  accumulation, morphologic signs of mitochondrial swelling (Puka-Sundvall et al., 2000a), and an increase in mitochondrial entrapment of  $^{14}\text{C}$ -labeled 2-deoxyglucose shortly after hypoxic ischemia (Puka-Sundvall et al., 2001). Although significant evidence suggests involvement of the PTP in adult animal models of TBI (Sullivan et al., 2005), this has not been directly investigated in pediatric models. Our finding of early mitochondrial release of cytochrome *c* after TBI in immature rats could be secondary to mitochondrial swelling via PTP opening, but this could also occur from other mechanisms. Future studies examining mitochondrial swelling and potential neuroprotection with CsA in immature animal models of TBI would be informative.

#### Mitochondria and the developing brain

Developmental differences in brain mitochondria of normal rats have been well documented (Table 1). In general, through the first 3–4 weeks of life in the rat, there is a 3-fold increase in mitochondrial protein per cell, with corresponding increases in respiratory enzyme activity and increasing oxygen consumption (Tyler and Harrevel, 2004).

**Table 1**

Unique age-dependent features of cellular response to acute brain injury from immature animal studies.

Mechanism	Unique response of developing brain	References
Energy metabolism	<ul style="list-style-type: none"> <li>Improved potential for use of alternative fuels (ketones)</li> </ul>	(Thomas et al., 2000; Prins et al., 2005)
Excitotoxicity	<ul style="list-style-type: none"> <li>Shortened length of metabolic depression after TBI</li> <li>Regional expression of NMDA, AMPA and kainate receptors is age-dependent</li> <li>Heightened sensitivity of young brain to excitotoxicity after HI insult</li> <li>Potential for NMDA inhibition to promote delayed (apoptotic) cell death</li> <li>Potential for GABA receptor excitatory transmission at young ages</li> <li>Age-dependent differences in mitochondrial calcium uptake capacity</li> </ul>	(McDonald et al., 1988; McDonald et al., 1992; Monyer et al., 1994; Ben Ari et al., 1997; Pohl et al., 1999; Anderson et al., 1999; Sharma et al., 2000; Robertson et al., 2004)
Mitochondrial permeability transition	<ul style="list-style-type: none"> <li>Lack of effect of cyclosporin A (CsA) on mitochondrial <math>\text{Ca}^{2+}</math> uptake or HI injury</li> <li>CsA protection with intrauterine ischemia</li> <li>CsA worsened mortality (PND 6) and seizure length (PND 30) in immature rats</li> </ul>	(Puka-Sundvall et al., 2001; Robertson et al., 2004; Nakai et al., 2004; Setkowicz et al., 2004)
Oxidative stress	<ul style="list-style-type: none"> <li>Brain antioxidant enzymes have an age-dependent profile of activity and expression</li> <li>Perinatal surges in antioxidant enzymes but lower levels of enzymes during other developmental times (versus adult)</li> <li>Worse outcome after HI insult in Cu,Zn-SOD overexpressing immature mice (PND 7) due to accumulation of <math>\text{H}_2\text{O}_2</math></li> <li>Limited upregulation of GPX after TBI in immature mouse (PND 21) compared to adult mouse</li> </ul>	(Mavelli et al., 1982; Ditelberg et al., 1996; Fullerton et al., 1998; Khan and Black, 2003; Fan et al., 2003; Robertson et al., 2006; Bayir et al., 2006)
Cell death	<ul style="list-style-type: none"> <li>Immature brain has normal, basal programmed cell death mechanisms active in early development</li> <li>Age-dependent expression of bcl-2 family proteins in tissue and mitochondria</li> <li>Susceptibility of immature brain to apoptotic cell death after hypoxic-ischemic insult</li> <li>Vulnerability of young brain to mitochondrial cytochrome <i>c</i> release upon exposure to pro-apoptotic stimuli</li> <li>Trauma-induced apoptosis is greater in young rats</li> </ul>	(Raff et al., 1993; Alonso et al., 1997; Shimohama et al., 1998; Bittigau et al., 1999; Krajewska et al., 2002; Johnston et al., 2002; Polster et al., 2003b; Vannucci and Hagberg, 2004; Zhu et al., 2005; Northington et al., 2005; Soane et al., 2008)

TBI – traumatic brain injury; HI – hypoxic-ischemic; PND – postnatal day; SOD – superoxide dismutase; GPX – glutathione peroxidase.

1942; Murthy and Rappoport, 1963; Milstein et al., 1968). There are also potential differences in mitochondrial membrane composition (Sitkiewicz et al., 1982), and relative ratios of synaptosomal to non-synaptosomal brain mitochondria (Dienel et al., 1977). A series of studies by the laboratory of Holtzman and others have detailed developmental differences in brain mitochondrial activity in immature (<4 weeks old) versus mature rats (adult, >2 months). ADP/O ratios with NAD-lined substrates were lower in rats <2 weeks of age, increased between the 3rd and 4th week, and reached adult levels by the 4th week of life (Holtzman and Moore, 1973; Holtzman and Moore, 1975). There are also differences in State III and State IV respiratory rates at different ages, reaching adult levels at >5 weeks of age (Holtzman and Moore, 1975). Additional studies of mitochondrial function demonstrate increased sensitivity of cortical slices from immature animals to altered pH, with a lower pH (6.7) inhibiting immature tissue respiration but not affecting adult tissue respiration (Holtzman et al., 1987).

Developmental aspects of mitochondrial susceptibility to brain insults have been shown using models of hypoxia–ischemia in immature rats (7 day old). A comparative *in vitro* study of isolated nerve terminals after hypoxic ischemia demonstrated differences in mitochondrial complex activity in the neonatal versus adult tissues (Keelan et al., 1999). *In vivo* studies of hypoxic ischemia in immature rats (7 day old) have shown significant abnormalities in mitochondrial respiration patterns (Yager et al., 1991a; Yager et al., 1991b; Yager et al., 1996; Gilland et al., 1998). The apoptotic pathway also has a role in neurodegeneration following hypoxic ischemia in the developing rat brain. In 21 day old rats, evidence of apoptosis was seen after both mild and severe unilateral hypoxic ischemic injuries (Beilharz et al., 1995). Younger rats (7 day old) had evidence of apoptotic changes throughout the brain, with prolonged increases in caspase-3 activity (Nakajima et al., 2000; Puka-Sundvall et al., 2000b; Puka-Sundvall et al., 2000c; Puka-Sundvall et al., 2000d). This unique pattern of activated caspase-3 was also seen after hypoxic ischemia in rats of various ages, with the youngest rats (7 day old) having the highest levels of active caspase-3, that declined steadily during maturation (Hu et al., 2000). Perinatal models have shown alteration in mitochondrial mRNA and DNA levels associated with uteroplacental insufficiency in rats (Lane et al., 2000), and histologic evidence of apoptotic cell morphology on brain autopsy tissue from infants suffering intrauterine cerebral injury (Edwards et al., 1997). This apparent propensity toward apoptosis could be explained at least in part by a greater general expression of pro-apoptotic compared to anti-apoptotic proteins at the level of brain mitochondria and the greater sensitivity to release of mitochondrial apoptotic mediators, e.g., cytochrome c, in response to apoptotic triggers, including BH3 only proteins and elevated intracellular  $Ca^{2+}$ , as demonstrated by our lab and the Hagberg group (Polster et al., 2003c; Zhu et al., 2005; Soane et al., 2008).


Studies in pediatric TBI models have also demonstrated mitochondrial dysfunction and altered cerebral metabolism (Robertson et al., 2006). Investigators have found early hyperglycolysis followed by a 1–3 day period of metabolic depression in fluid percussion TBI in immature rats (17 day old) (Thomas et al., 2000). This group has also shown that the ability of alternative substrates, such as ketone bodies, to rescue brain metabolism may be developmentally regulated (Prins et al., 2005). Using the controlled cortical impact model, we studied isolated brain mitochondria early (1–4 h) after TBI in immature rats (17 day old) (Robertson et al., 2007). Mitochondria had significant alterations in respiratory capacity, with increases in State 4 respiration (1 h) and decreases in State 3 respiration (4 h). Mitochondria also had reduced cytochrome c content and decreased activity of the important metabolic enzyme, pyruvate dehydrogenase. Other investigators used gel-based proteomics to show significant reductions in pyruvate dehydrogenase subunit expression that extended to 2 weeks after TBI in the

developing brain (Kochanek et al., 2006). Loss of pyruvate dehydrogenase activity in pediatric TBI may be especially important, as it is the critical enzymatic link between glycolysis and the TCA cycle. Evidence for metabolic derangements in clinical studies of pediatric TBI comes primarily from imaging investigations (Ashwal et al., 2006). Using magnetic resonance spectroscopy (MRS), children have marked elevations in brain lactate and reductions in N-acetyl aspartate (NAA), a marker of neuronal and/or mitochondrial integrity, and these metabolic alterations correlate with long-term neurologic outcome (Ashwal et al., 2000). However, few of these imaging studies were obtained early after pediatric TBI (<48 h), so information is not available about very early metabolic alterations. Recently, we used proton ( $^1H$ ) spectroscopy after CCI in immature rats (16–17 day old) to evaluate the time course of metabolic alterations (Casey et al., 2008). We showed that metabolic derangements begin early (<4 h) and are sustained for at least 7 days after TBI in the developing brain. Another study used  $^{13}C$ -NMR spectroscopy to evaluate glucose metabolism 5–6 h after CCI in immature rats (21–22 day old) (Scafidi et al., 2009). This study showed that neuronal oxidative metabolism of glucose is delayed in both the ipsilateral and contralateral hemispheres, compared to uninjured sham controls. Specifically, findings suggested impairment in the TCA cycle, at the level of pyruvate dehydrogenase and possibly  $\alpha$ -ketoglutarate dehydrogenase, as well as impairment of the malate–aspartate shuttle. Future studies using clinically relevant tools, such as MR spectroscopy, could allow development and testing of mitochondrially targeted neuroprotective strategies for the developing brain after injury. This study demonstrates the value of utilizing clinically relevant tools, such as MRS, to investigate metabolic alterations following acute pediatric brain injury.

#### *Mitochondrial pathways for neuroprotection in the immature brain*

Based on increasing knowledge of mitochondrial pathways toward cell death in the immature brain after hypoxia, ischemia, and trauma, several mitochondria-based neuroprotective strategies have been developed (Fig. 2). One approach is pharmacologic inhibition of mitochondrial  $Ca^{2+}$  overload, which together with oxidative stress, induces opening of the inner membrane permeability transition pore. While cyclosporin-based drugs appear effective at inhibiting PTP opening in mature brain mitochondria, the relatively low expression of cyclophilin D in immature brain mitochondria may be responsible for the ineffectiveness of cyclosporin A at improving outcome after neonatal brain injury (Puka-Sundvall et al., 2001; Eliseev et al., 2007). Other attempts at PTP inhibition should be made, however, including the use of agents, e.g., 2-aminoethoxydiphenyl borate, that are capable of inhibiting cyclosporin A-insensitive PTP (Chinopoulos et al., 2003). Another approach toward PTP inhibition includes the use of agonists for mitochondrial ATP-sensitive potassium channels, that indirectly inhibit PTP opening in adult brain mitochondria (Wu et al., 2006). These agents, e.g., diazoxide, also inhibit mitochondrial ROS production (Fornazari et al., 2008) and have exhibited neuroprotection in a neonatal hypoxic ischemia model (Rajapakse et al., 2002).

One consequence of PTP opening is loss of mitochondrial pyridine nucleotides, that are required in numerous reactions involved in normal energy metabolism and in ROS detoxification. Therapies directed at replenishing cellular and mitochondrial cofactors, e.g., NAD (H), have exhibited neuroprotection both *in vitro* and with adult animal models of brain injury and should be tested with pediatric brain injury models. This strategy includes administration of  $NAD^+$  or nicotinamide (Ayoub et al., 1999; Sadanaga-Akiyoshi et al., 2003; Ying et al., 2007). One report indicated that when nicotinamide was administered to neonatal rats as long as 24 h after perinatal asphyxia, dopamine levels present within the striatum 3 months later were relatively preserved (Bustamante et al., 2007). Moreover, since a futile



**Fig. 2.** Mitochondria-targeted neuroprotective interventions for acute injury to the immature brain. One common early trigger of neuronal injury following either cerebral ischemia or head trauma is an abnormal elevation of intracellular  $\text{Ca}^{2+}$ , which if unabated, will directly lead to cellular necrosis through activation of degradative enzymes, e.g., phospholipases and proteases. Elevated  $\text{Ca}^{2+}$  can also induce delayed cell death by inducing mitochondrial metabolic dysfunction, e.g., through activation of the permeability transition pore (PTP), and by stimulation of activating enzymes that produce reactive O<sub>2</sub> species (ROS), e.g., NADPH oxidase (NOX) and nitric oxide synthase (NOS). The ROS contribute to mitochondrial metabolic dysfunction, stimulate cellular inflammatory responses, e.g., microglial activation, and modify many different molecules throughout the cell including lipids, proteins, and DNA. DNA damage activates poly-ADP-ribose polymerase, which consumes  $\text{NAD}^+$ , thus impairing energy metabolism. Pro-apoptotic molecular pathways, including those activated by inflammatory cytokines, lead to the release of mitochondrial apoptotic proteins, e.g., cytochrome *c* (Cyt *c*) and apoptosis-inducing factor (AIF). Calpain cleavage of molecules, e.g., AIF and Bid, also promotes the mitochondrial intrinsic pathway of apoptosis. Potentially neuroprotective interventions that target these pathways include the PTP inhibitor 2-aminoethoxydiphenyl borate (2-APB), the ATP-sensitive  $\text{K}^+$  channel opener diazoxide, normoxic resuscitation, the Nrf2 activator sulforaphane (SFP), calpain inhibitors, the PARP inhibitor 3-amino benzamide (3ABA), estrogen, which induces Bcl2 expression, Bax inhibitors, and several agents that stimulate aerobic energy metabolism, including free carnitine and acetyl-L-carnitine,  $\beta$ -hydroxybutyrate ( $\beta\text{OHB}$ ),  $\text{NAD}^+$  or its precursor nicotinamide, and pyruvate.

cycle of  $\text{NAD}^+$  catabolism is generated by the activation of both PARP and PARG, inhibitors of these enzymes should preserve both cytosolic and mitochondrial energy metabolism. Poly-ADP ribosylation of proteins occurs both early (<1 h) and at delayed times (12 h) after stroke in neonates (Martin et al., 2005). The PARP inhibitor 3-aminobenzamide both inhibits this product formation and subsequent, delayed cell death (Joly et al., 2003). Although not yet reported for immature animals, TBI in adult rats results in mitochondrial protein poly-ADP-ribosylation that may contribute to inhibition of respiration and oxidative phosphorylation (Lai et al., 2008). Additional investigation of this form of metabolic impairment in the immature brain is clearly needed.

Another mitochondria-based approach to neuroprotection for acute brain injury is the administration of agents that in one way or another act as enhancers of aerobic energy metabolism. The administration of exogenous pyridine nucleotides and their precursors, as mentioned above, is one example. Other examples include agents that can directly serve as fuel for cerebral energy metabolism, e.g., ketone bodies, acetyl-L-carnitine, and pyruvate. Of these,  $\beta$ -hydroxybutyrate may be the most straightforward approach for the immature brain since the vascular monocarboxylic acid transporters that carry ketone bodies and the brain enzymes that metabolize them are relatively high during the suckling period (Vannucci and Simpson, 2003). Thus, the immature brain is genomically primed to utilize ketones as fuel, particularly if the activity of enzymes, e.g., pyruvate dehydrogenase, become inhibited, limiting aerobic glucose metabolism.  $\beta$ -hydroxybutyrate, a ketogenic

diet, or fasting has been shown to be neuroprotective in various rat models of acute brain injury (Yamada et al., 2005; Puchowicz et al., 2008; Tai et al., 2008; Davis et al., 2008), with at least one report for young rats (Prins et al., 2005). While a ketogenic diet is used to treat intractable epilepsy in children, there is some concern that it can also impair normal brain development (Zhao et al., 2004). An alternative approach, based on the same principle of compensating for impaired glucose metabolism, is administration of acetyl-L-carnitine or free carnitine. These natural agents normally involved in acetyl and acyl carbon metabolism are neuroprotective and promote the production of acetyl-CoA, the same product generated from glucose derived pyruvate via pyruvate dehydrogenase. Large doses of L-carnitine administered prior to, but not after neonatal hypoxic ischemia in rats result in neuroprotection (Wainwright et al., 2003). While carnitine appears to improve cerebral energy metabolism after hypoxic ischemia (Macri et al., 1995), it may exert mechanisms of action independent of directly promoting mitochondrial energy metabolism (Winter et al., 1995; Akisu et al., 1998; Tastekin et al., 2005; Wainwright et al., 2006). Acetyl-L-carnitine has not been tested for neuroprotection with immature animals; however, it is highly effective against brain injury in adult animals (Rosenthal et al., 1992; Shuaib et al., 1995; Lolic et al., 1997; Zanelli et al., 2005).

Pyruvate is a third metabolic substrate that has exhibited neuroprotection in different acute brain injury models; however, it has not yet been reported using immature animals. The brain mitochondrial redox state is relatively oxidized soon after neonatal hypoxic ischemia (Yager et al., 1991a), which could be explained by either a limited concentration of pyruvate generated by glycolysis or by limited influx of carbon into the tricarboxylic acid cycle through impaired pyruvate dehydrogenase. If the first scenario is true, exogenous pyruvate should promote aerobic energy metabolism, possibly resulting in neuroprotection. The same result might even be reached by adding lactate, as endogenous lactate is a preferred substrate for immature brain metabolism, at least during anoxia and reoxygenation (Vannucci and Duffy, 1976). Stimulation of metabolism through pyruvate dehydrogenase can also occur by addition of dichloroacetate, which inhibits pyruvate dehydrogenase kinase, thereby de-inhibiting pyruvate dehydrogenase (Stacpoole, 1989). Dichloroacetate has exhibited protection for both ischemia and trauma in the adult brain (Chang et al., 1992; Biros and Nordness, 1996). In addition, one study reported that early post-treatment with dichloroacetate improved post-ischemic clearance of cerebral acidosis and elevated phosphocreatine in newborn pigs better than it did for 1 month old swine (Corbett et al., 1998); however, effects on neurologic or histologic outcome were not reported.

A host of other interventions could prove therapeutic for acute injury to the immature brain that indirectly compensate for mitochondrial dysfunction or that protect against mitochondrial damage from conditions, e.g., elevated intracellular  $\text{Ca}^{2+}$  and high levels of ROS, that exist during and after ischemia and trauma. For instance, pretreatment of rats with creatine was reported to improve the cerebral phosphocreatine/phosphate ratio and reduce brain edema after neonatal hypoxic ischemia (Adcock et al., 2002). Elevating total brain creatine may provide neuroprotection by elevating the high energy phosphate buffering capacity through accelerating the mitochondrial creatine kinase reaction. N-acetylcysteine, an antioxidant, was also shown to reduce cortical  $\text{H}_2\text{O}_2$  and brain lactate and improve cerebral perfusion in piglets after hypoxia, suggesting an improvement in cerebral aerobic energy metabolism (Lee et al., 2008). In addition to avoiding brain tissue hyperoxia and adding antioxidants to reduce oxidative stress, a new approach has been reported that is based on activation of endogenous antioxidant and anti-inflammatory gene expression. Low levels of ROS and specific chemicals, e.g. sulforaphane, an isothiocyanate found in cruciferous vegetables, promote the translocation of Nrf2, a transcriptional activating factor, from the cytosol to the nucleus, where it binds to antioxidant response elements (ARE), thereby

stimulating the expression of scores of genes (Jaiswal, 2004; Dinkova-Kostova and Talalay, 2008). Some of the elevated gene products appear in mitochondria, including Mn-superoxide dismutase, glutathione peroxidase, and glutathione-S-transferase (Zhu et al., 2008). Considering that the immature brain has relatively low expression levels for antioxidant genes (Cruz et al., 1998; Bayir et al., 2006), an antioxidant genomic post-conditioning strategy could be particularly effective toward neuroprotection in infants and children.

Since neuronal apoptosis appears even more prominent following acute injury to the immature brain than it does in adults, interventions that target the mitochondrial (intrinsic) pathway of apoptosis are very promising. For instance, the release of apoptosis-inducing factor (AIF) and its role in DNA damage is important in neonatal hypoxic ischemic brain injury (Zhu et al., 2003). The Ca<sup>2+</sup> activated protease calpain plays a critical role in mitochondrial AIF release (Polster et al., 2005; Cao et al., 2007), and one report indicates that calpain inhibitors are neuroprotective in neonatal rat cerebral hypoxic ischemia (Chen et al., 2008). Other anti-apoptotic approaches that could be applied to the immature brain include the use of agents, e.g., estrogen, that stimulate the expression of Bcl-2 and other apoptotic proteins (Zhang et al., 2004), or agents that impair the ability of Bax or Bad to create pores in the mitochondrial inner membrane resulting in the release of cytochrome *c* and other pro-apoptotic proteins into the cytosol (Polster et al., 2003a; Guan et al., 2006; Iriyama et al., 2008).

## Acknowledgments

These authors were supported by NIH grants K08 NS43805, R01 NS34152, and P01 HD16596.

## References

- Adcock, K.H., Nedelcu, J., Loenneker, T., Martin, E., Wallimann, T., Wagner, B.P., 2002. Neuroprotection of creatine supplementation in neonatal rats with transient cerebral hypoxia-ischemia. *Dev. Neurosci.* 24, 382–388.
- Adelson, P.D., Bratton, S.L., Carney, N.A., Chesnut, R.M., du Coudray, H.E., Goldstein, B., Kochanek, P.M., Miller, H.C., Partington, M.D., Selden, N.R., Warden, C.W., Wright, D.W., 2003. Guidelines for the acute medical management of severe traumatic brain injury in infants, children, and adolescents. Chapter 17. Critical pathway for the treatment of established intracranial hypertension in pediatric traumatic brain injury. *Pediatr. Crit. Care Med.* 4, S65–S67.
- Adelson, P.D., Ragheb, J., Kanev, P., Brockmeyer, D., Beers, S.R., Brown, S.D., Cassidy, L.D., Chang, Y., Levin, H., 2005. Phase II clinical trial of moderate hypothermia after severe traumatic brain injury in children. *Neurosurgery* 56, 740–754.
- Ahn, E.S., Robertson, C.L., Vereczki, V., Hoffman, G.E., Fiskum, G., 2008. Normoxic ventilatory resuscitation following controlled cortical impact reduces peroxynitrite-mediated protein nitration in the hippocampus. *J. Neurosurg.* 108, 124–131.
- Akisu, M., Kultursay, N., Coker, I., Huseynov, A., 1998. The effect of L-carnitine on platelet activating factor concentration in the immature rat model of hypoxic-ischemic brain injury. *Acta Med. Okayama.* 52, 183–187.
- Alessandri, B., Rice, A.C., Levasseur, J., DeFord, M., Hamm, R.J., Bullock, M.R., 2002. Cyclosporin A improves brain tissue oxygen consumption and learning/memory performance after lateral fluid percussion injury in rats. *J. Neurotrauma* 19 (7), 829–841.
- Alonso, G., Guillemain, I., Dumoulin, A., Privat, A., Patey, G., 1997. Immunolocalization of Bcl-xL/S in the central nervous system of neonatal and adult rats. *Cell. Tissue Res.* 288, 59–68.
- Anderson, K.J., Mason, K.L., McGraw, T.S., Theophilopoulos, D.T., Sapper, M.S., Burchfield, D.J., 1999. The ontogeny of glutamate receptors and D-aspartate binding sites in the ovine CNS. *Brain Res. Dev. Brain Res.* 118, 69–77.
- Anderson, V., Catroppa, C., Morse, S., Haritou, F., Rosenfeld, J., 2005. Functional plasticity or vulnerability after early brain injury? *Pediatrics* 116, 1374–1382.
- Ashwal, S., Holshouser, B.A., Shu, S.K., Simmons, P.L., Perkin, R.M., Tomasi, L.G., Knierim, D.S., Sheridan, C., Craig, K., Andrews, G.H., Hinshaw, D.B., 2000. Predictive value of proton magnetic resonance spectroscopy in pediatric closed head injury. *Pediatr. Neurol.* 23, 114–125.
- Ashwal, S., Holshouser, B.A., Tong, K.A., 2006. Use of advanced neuroimaging techniques in the evaluation of pediatric traumatic brain injury. *Dev. Neurosci.* 28, 309–326.
- Ayoub, I.A., Lee, E.J., Ogilvy, C.S., Beal, M.F., Maynard, K.I., 1999. Nicotinamide reduces infarction up to two hours after the onset of permanent focal cerebral ischemia in Wistar rats. *Neurosci. Lett.* 259, 21–24.
- Azzopardi, D., Brocklehurst, P., Edwards, D., Halliday, H., Levene, M., Thoresen, M., Whitelaw, A., 2008. The TOBY Study. Whole body hypothermia for the treatment of perinatal asphyxial encephalopathy: a randomised controlled trial. *BMC Pediatr.* 8, 17.
- Balan, I.S., Fiskum, G., Hazelton, J., Cotto-Cumba, C., Rosenthal, R.E., 2006. Oximetry-guided reoxygenation improves neurological outcome after experimental cardiac arrest. *Stroke* 37 (12), 3008–3013.
- Barton, C., Callahan, M., 1991. High-dose epinephrine improves the return of spontaneous circulation rates in human victims of cardiac arrest. *Ann. Emerg. Med.* 20 (7), 722–725.
- Bayir, H., Kochanek, P.M., Kagan, V.E., 2006. Oxidative stress in immature brain after traumatic brain injury. *Dev. Neurosci.* 28, 420–431.
- Beilharz, E.J., Williams, C.E., Dragunow, M., Sirimanne, E.S., Gluckman, P.D., 1995. Mechanisms of delayed cell death following hypoxic-ischemic injury in the immature rat: evidence for apoptosis during selective neuronal loss. *Brain Res. Mol. Brain Res.* 29, 1–14.
- Ben Ari, Y., Khazipov, R., Leinekugel, X., Caillard, O., Gaiarsa, J.L., 1997. GABAA, NMDA and AMPA receptors: a developmentally regulated 'menage a trois'. *Trends Neurosci.* 20, 523–529.
- Berg, M.D., Nadkarni, V.M., Berg, R.A., 2008. Cardiopulmonary resuscitation in children. *Curr. Opin. Crit. Care.* 14, 254–260.
- Bernard, S.A., Gray, T.W., Buist, M.D., Jones, B.M., Silvester, W., Gutteridge, G., Smith, K., 2002. Treatment of comatose survivors of out-of-hospital cardiac arrest with induced hypothermia. *N. Engl. J. Med.* 346, 557–563.
- Bernardi, P., Scorrano, L., Colonna, R., Petronilli, V., Di Lisa, F., 1999. Mitochondria and cell death. Mechanistic aspects and methodological issues. *Eur. J. Biochem.* 264 (3), 687–701.
- Biros, M.H., Nordness, R., 1996. Effects of chemical pretreatment on posttraumatic cortical edema in the rat. *Am. J. Emerg. Med.* 14, 27–32.
- Biswas, A.K., Bruce, D.A., Sklar, F.H., Bokovoy, J.L., Sommerauer, J.F., 2002. Treatment of acute traumatic brain injury in children with moderate hypothermia improves intracranial hypertension. *Crit. Care Med.* 30, 2742–2751.
- Bittigau, P., Siffringer, M., Pohl, D., Stadthaus, D., Ishimaru, M., Shimizu, H., Ikeda, M., Lang, D., Speer, A., Olney, J.W., Ikonomidou, C., 1999. Apoptotic neurodegeneration following trauma is markedly enhanced in the immature brain. *Ann. Neurol.* 45, 724–735.
- Blomgren, K., Hagberg, H., 2006. Free radicals, mitochondria, and hypoxia-ischemia in the developing brain. *Free Radic. Biol. Med.* 40 (3), 388–397.
- Bogaert, Y.E., Rosenthal, R.E., Fiskum, G., 1994. Posts ischemic inhibition of cerebral cortex pyruvate dehydrogenase. *Free Radic. Biol. Med.* 16 (6), 811–820.
- Brain Resuscitation Clinical Trial I Study Group, 1986. Randomized clinical study of thiopental loading in comatose survivors of cardiac arrest. *N. Engl. J. Med.* 314, 397–403.
- Brain Resuscitation Clinical Trial II Study Group, 1991. A randomized clinical study of a calcium-entry blocker (lidoflazine) in the treatment of comatose survivors of cardiac arrest. *N. Engl. J. Med.* 324, 1225–1231.
- Broccard, A., 2006. Therapeutic hypothermia for anoxic brain injury following cardiac arrest: a "cool" transition toward cardiopulmonary cerebral resuscitation. *Crit. Care Med.* 34 (7), 2008–2009.
- Bustamante, D., Morales, P., Pereyra, J.T., Gojny, M., Herrera-Marschitz, M., 2007. Nicotinamide prevents the effect of perinatal asphyxia on dopamine release evaluated with in vivo microdialysis 3 months after birth. *Exp. Brain Res.* 177, 358–369.
- Cao, G., Xing, J., Xiao, X., Liou, A.K., Gao, Y., Yin, X.M., Clark, R.S., Graham, S.H., Chen, J., 2007. Critical role of calpain I in mitochondrial release of apoptosis-inducing factor in ischemic neuronal injury. *J. Neurosci.* 27, 9278–9293.
- Casey, P.A., McKenna, M.C., Fiskum, G., Saraswati, M., Robertson, C.L., 2008. Early and sustained alterations in cerebral metabolism after traumatic brain injury in immature rats. *J. Neurotrauma* 25, 603–614.
- Castino, R., Bellio, N., Nicotra, G., Follo, C., Trincheri, N.F., Isidoro, C., 2007. Cathepsin D-Bax death pathway in oxidative stressed neuroblastoma cells. *Free Radic. Biol. Med.* 42 (9), 1305–1316.
- Chang, L.H., Shimizu, H., Abiko, H., Swanson, R.A., Faden, A.I., James, T.L., Weinstein, P.R., 1992. Effect of dichloroacetate on recovery of brain lactate, phosphorus energy metabolites, and glutamate during reperfusion after complete cerebral ischemia in rats. *J. Cereb. Blood Flow Metab.* 12, 1030–1038.
- Chen, L.N., Yan, B., Chen, D.P., Yao, Y.J., 2008. [Protective effect of calpain inhibitor-3 on hypoxic-ischemic brain damage of neonatal rats. *Zhonghua Er Ke Za Zhi* 46, 13–17.
- Chinopoulos, C., Starkov, A.A., Fiskum, G., 2003. Cyclosporin A-insensitive permeability transition in brain mitochondria: inhibition by 2-aminoethoxydiphenyl borate. *J. Biol. Chem.* 278, 27382–27389.
- Contreras, L., Satrustegui, J., 2009. Calcium signaling in brain mitochondria: interplay of malate aspartate NADH shuttle and calcium uniporter/mitochondrial dehydrogenase pathways. *J. Biol. Chem.* 284, 7091–7099.
- Corbett, R., Lupton, A., Gee, J., Garcia, D., Silmon, S., Tollefsbol, G., 1998. Age-related differences in the effect of dichloroacetate on posts ischemic lactate and acid clearance measured in vivo using magnetic resonance spectroscopy and microdialysis. *J. Neurochem.* 71, 1205–1214.
- Crompton, M., Barksby, E., Johnson, N., Capano, M., 2002. Mitochondrial intermembrane junctional complexes and their involvement in cell death. *Biochimie* 84 (2–3), 143–152.
- Cruz, F., Scott, S.R., Barroso, I., Santesteban, P., Cerdan, S., 1998. Ontogeny and cellular localization of the pyruvate recycling system in rat brain. *J. Neurochem.* 70, 2613–2619.
- Davis, L.M., Pauly, J.R., Readnower, R.D., Rho, J.M., Sullivan, P.G., 2008. Fasting is neuroprotective following traumatic brain injury. *J. Neurosci. Res.* 86, 1812–1822.
- Dienel, G., Ryder, E., Greengard, O., 1977. Distribution of mitochondrial enzymes between the perikaryal and synaptic fractions of immature and adult rat brain. *Biochim. Biophys. Acta* 496, 484–494.
- Dinkova-Kostova, A.T., Talalay, P., (2008) Direct and indirect antioxidant properties of inducers of cytoprotective proteins. *Mol. Nutr. Food Res.* 52 Suppl. 1:S128–38.: S128–S138.

- Diringer, M.N., 2008. Hyperoxia: good or bad for the injured brain? *Curr. Opin. Crit. Care* 14, 167–171.
- Ditelberg, J.S., Sheldon, R.A., Epstein, C.J., Ferriero, D.M., 1996. Brain injury after perinatal hypoxia-ischemia is exacerbated in copper/zinc superoxide dismutase transgenic mice. *Pediatr. Res.* 39, 204–208.
- Donoghue, A.J., Nadkarni, V., Berg, R.A., Osmond, M.H., Wells, G., Nesbitt, L., Stiell, I.G., 2005. Out-of-hospital pediatric cardiac arrest: an epidemiologic review and assessment of current knowledge. *Ann. Emerg. Med.* 46, 512–522.
- Du, L., Zhang, X., Han, Y.Y., Burke, N.A., Kochanek, P.M., Watkins, S.C., Graham, S.H., Carcillo, J.A., Szabo, C., Clark, R.S., 2003. Intra-mitochondrial poly(ADP-riboseylation) contributes to NAD<sup>+</sup> depletion and cell death induced by oxidative stress. *J. Biol. Chem.* 278 (20), 18426–18433.
- Edwards, A.D., Yue, X., Cox, P., Hope, P.L., Azzopardi, D.V., Squier, M.V., Mehmet, H., 1997. Apoptosis in the brains of infants suffering intrauterine cerebral injury. *Pediatr. Res.* 42, 684–689.
- Eliasson, M.J., Sampei, K., Mandir, A.S., Hurn, P.D., Traystman, R.J., Bao, J., Pieper, A., Wang, Z.Q., Dawson, T.M., Snyder, S.H., Dawson, V.L., 1997. Poly(ADP-ribose) polymerase gene disruption renders mice resistant to cerebral ischemia. *Nat. Med.* 3 (10), 1089–1095.
- Eliseev, R.A., Filippov, G., Velos, J., Vanwinkle, B., Goldman, A., Rosier, R.N., Gunter, T.E., 2007. Role of cyclophilin D in the resistance of brain mitochondria to the permeability transition. *Neurobiol. Aging* 28, 1532–1542.
- Ewing-Cobbs, L., Barnes, M., Fletcher, J.M., Levin, H.S., Swank, P.R., Song, J., 2004a. Modeling of longitudinal academic achievement scores after pediatric traumatic brain injury. *Dev. Neuropsychol.* 25, 107–133.
- Ewing-Cobbs, L., Prasad, M.R., Kramer, L., Cox Jr., C.S., Baumgartner, J., Fletcher, S., Mendez, D., Barnes, M., Zhang, X., Swank, P., 2006. Late intellectual and academic outcomes following traumatic brain injury sustained during early childhood. *J. Neurosurg.* 105, 287–296.
- Ewing-Cobbs, L., Prasad, M.R., Landry, S.H., Kramer, L., DeLeon, R., 2004b. Executive functions following traumatic brain injury in young children: a preliminary analysis. *Dev. Neuropsychol.* 26, 487–512.
- Fabian, R.H., Perez-Polo, J.R., Kent, T.A., 2008. Perivascular nitric oxide and superoxide in neonatal cerebral hypoxia-ischemia. *Am. J. Physiol. Heart Circ. Physiol.* 295, H1809–H1814.
- Fan, P., Yamauchi, T., Noble, L.J., Ferriero, D.M., 2003. Age-dependent differences in glutathione peroxidase activity after traumatic brain injury. *J. Neurotrauma* 20, 437–445.
- Fiskum, G., Murphy, A.N., Beal, M.F., 1999. Mitochondria in neurodegeneration: acute ischemia and chronic neurodegenerative diseases. *J. Cereb. Blood Flow Metab.* 19 (4), 351–369.
- Fornazari, M., de Paula, J.G., Castilho, R.F., Kowaltowski, A.J., 2008. Redox properties of the adenosine triphosphate-sensitive K<sup>+</sup> channel in brain mitochondria. *J. Neurosci. Res.* 86, 1548–1556.
- Forte, M., Gold, B.G., Marracci, G., Chaudhary, P., Basso, E., Johnsen, D., Yu, X., Fowlkes, J., Rahder, M., Stem, K., Bernardi, P., Bourdette, D., 2007. Cyclophilin D inactivation protects axons in experimental autoimmune encephalomyelitis, an animal model of multiple sclerosis. *Proc. Natl. Acad. Sci. U.S.A.* 104 (18), 7558–7563.
- Fullerton, H.J., Ditelberg, J.S., Chen, S.F., Sarco, D.P., Chan, P.H., Epstein, C.J., Ferriero, D.M., 1998. Copper/zinc superoxide dismutase transgenic brain accumulates hydrogen peroxide after perinatal hypoxia ischemia. *Ann. Neurol.* 44, 357–364.
- Gilland, E., Puka-Sundvall, M., Hillered, L., Hagberg, H., 1998. Mitochondrial function and energy metabolism after hypoxia-ischemia in the immature rat brain: involvement of NMDA-receptors. *J. Cereb. Blood Flow Metab.* 18, 297–304.
- Gluckman, P.D., Wyatt, J.S., Azzopardi, D., Ballard, R., Edwards, A.D., Ferriero, D.M., Polin, R.A., Robertson, C.M., Thoresen, M., Whitelaw, A., Gunn, A.J., 2005. Selective head cooling with mild systemic hypothermia after neonatal encephalopathy: multicentre randomised trial. *Lancet* 365, 663–670.
- Goetting, M.G., Paradis, N.A., 1991. High-dose epinephrine improves outcome from pediatric cardiac arrest. *Ann. Emerg. Med.* 20 (1), 22–26.
- Grafe, M.R., Woodworth, K.N., Noppens, K., Perez-Polo, J.R., 2008. Long-term histological outcome after post-hypoxic treatment with 100% or 40% oxygen in a model of perinatal hypoxic-ischemic brain injury. *Int. J. Dev. Neurosci.* 26, 119–124.
- Guan, Q.H., Pei, D.S., Liu, X.M., Wang, X.T., Xu, T.L., Zhang, G.Y., 2006. Neuroprotection against ischemic brain injury by SP600125 via suppressing the extrinsic and intrinsic pathways of apoptosis. *Brain Res.* 1092, 36–46.
- Gunn, A.J., Wyatt, J.S., Whitelaw, A., Barks, J., Azzopardi, D., Ballard, R., Edwards, A.D., Ferriero, D.M., Gluckman, P.D., Polin, R.A., Robertson, C.M., Thoresen, M., 2008. Therapeutic hypothermia changes the prognostic value of clinical evaluation of neonatal encephalopathy. *J. Pediatr.* 152 (55–8), 58.
- Halestrap, A.P., McStay, G.P., Clarke, S.J., 2002. The permeability transition pore complex: another view. *Biochimie* 84 (2–3), 153–166.
- Hansson, M.J., Mattiasson, G., Mansson, R., Karlsson, J., Keep, M.F., Waldmeier, P., Ruegg, U.T., Dumont, J.M., Besseghir, K., Elmer, E., 2004. The nonimmunosuppressive cyclosporin analogs NIM811 and UNIL025 display nanomolar potencies on permeability transition in brain-derived mitochondria. *J. Bioenerg. Biomembr.* 36 (4), 407–413.
- Harukuni, I., Bhardwaj, A., 2006. Mechanisms of brain injury after global cerebral ischemia. *Neurol. Clin.* 24 (1), 1–21.
- Hillered, L., Ernster, L., 1983. Respiratory activity of isolated rat brain mitochondria following in vitro exposure to oxygen radicals. *J. Cereb. Blood Flow Metab.* 3 (2), 207–214.
- Holtzman, D., Moore, C.L., 1973. Oxidative phosphorylation in immature rat brain mitochondria. *Biol. Neonate* 22, 230–242.
- Holtzman, D., Moore, C.L., 1975. Respiration in immature rat brain mitochondria. *J. Neurochem.* 24, 1011–1015.
- Holtzman, D., Olson, J.E., Nguyen, H., Hsu, J., Lewiston, N., 1987. Brain cellular and mitochondrial respiration in media of altered pH. *Metab. Brain Dis.* 2, 127–137.
- Hu, B.R., Liu, C.L., Ouyang, Y., Blomgren, K., Siesjo, B.K., 2000. Involvement of caspase-3 in cell death after hypoxia-ischemia declines during brain maturation. *J. Cereb. Blood Flow Metab.* 20, 1294–1300.
- Hutchison, J.S., Ward, R.E., Lacroix, J., Hebert, P.C., Barnes, M.A., Bohn, D.J., Dirks, P.B., Doucette, S., Fergusson, D., Gottesman, R., Joffe, A.R., Kirpalani, H.M., Meyer, P.G., Morris, K.P., Moher, D., Singh, R.N., Skippen, P.W., 2008. Hypothermia therapy after traumatic brain injury in children. *N. Engl. J. Med.* 358, 2447–2456.
- Hypothermia Cardiac Arrest Study Group, 2002. Mild therapeutic hypothermia to improve the neurologic outcome after cardiac arrest. *N. Engl. J. Med.* 346, 549–556.
- Iriyama, T., Kamei, Y., Kozuma, S., Taketani, Y., 2008. Bax-inhibiting peptide protects glutamate-induced cerebellar granule cell death by blocking Bax translocation. *Neurosci. Lett.* 451, 11–15.
- Jaiswal, A.K., 2004. Nrf2 signaling in coordinated activation of antioxidant gene expression. *Free Radic. Biol. Med.* 36, 1199–1207.
- Jankowitz, B.T., Adelson, P.D., 2006. Pediatric traumatic brain injury: past, present and future. *Dev. Neurosci.* 28, 264–275.
- Jastremski, M., Sutton-Tyrrell, K., Vaagenes, P., Abramson, N., Heiselman, D., Safar, P., 1989. Glucocorticoid treatment does not improve neurological recovery following cardiac arrest. *Brain Resuscitation Clinical Trial I Study Group. JAMA* 262, 3427–3430.
- Johnston, M.V., Nakajima, W., Hagberg, H., 2002. Mechanisms of hypoxic neurodegeneration in the developing brain. *Neuroscientist* 8, 212–220.
- Joly, L.M., Benjelloun, N., Plotkine, M., Charriat-Marlangue, C., 2003. Distribution of poly(ADP-ribose)ylation and cell death after cerebral ischemia in the neonatal rat. *Pediatr. Res.* 53, 776–782.
- Kauppinen, T.M., Swanson, R.A., 2007. The role of poly(ADP-ribose) polymerase-1 in CNS disease. *Neuroscience* 145 (4), 1267–1272.
- Keelan, J., Bates, T.E., Clark, J.B., 1999. Heightened resistance of the neonatal brain to ischemia-reperfusion involves a lack of mitochondrial damage in the nerve terminal. *Brain Res.* 821, 124–133.
- Keenan, H.T., Bratton, S.L., 2006. Epidemiology and outcomes of pediatric traumatic brain injury. *Dev. Neurosci.* 28, 256–263.
- Khan, J.Y., Black, S.M., 2003. Developmental changes in murine brain antioxidant enzymes. *Pediatr. Res.* 54, 77–82.
- Kochanek, A.R., Kline, A.E., Gao, W.M., Chadha, M., Lai, Y., Clark, R.S., Dixon, C.E., Jenkins, L.W., 2006. Gel-based hippocampal proteomic analysis 2 weeks following traumatic brain injury to immature rats using controlled cortical impact. *Dev. Neurosci.* 28, 410–419.
- Krajewska, M., Mai, J.K., Zapata, J.M., Ashwell, K.W., Schendel, S.L., Reed, J.C., Krajewski, S., 2002. Dynamics of expression of apoptosis-regulatory proteins Bid, Bcl-2, Bcl-X, Bax and Bak during development of murine nervous system. *Cell Death Differ.* 9, 145–157.
- Kuroda, S., Katsura, K.I., Tsuchida, R., Siesjo, B.K., 1996. Secondary bioenergetic failure after transient focal ischaemia is due to mitochondrial injury. *Acta Physiol. Scand.* 156 (2), 149–150.
- Lai, Y., Chen, Y., Watkins, S.C., Nathaniel, P.D., Guo, F., Kochanek, P.M., Jenkins, L.W., Szabo, C., Clark, R.S., 2008. Identification of poly-ADP-ribosylated mitochondrial proteins after traumatic brain injury. *J. Neurochem.* 104, 1700–1711.
- Lane, R.H., Tsirka, A.E., Gruetzmacher, E.M., 2000. Uteroplacental insufficiency alters cerebral mitochondrial gene expression and DNA in fetal and juvenile rats. *Pediatr. Res.* 47, 792–797.
- Lee, T.F., Tymafchuk, C.N., Bigam, D.L., Cheung, P.Y., 2008. Effects of postresuscitation N-acetylcysteine on cerebral free radical production and perfusion during reoxygenation of hypoxic newborn piglets. *Pediatr. Res.* 64, 256–261.
- Liu, Y., Rosenthal, R.E., Haywood, Y., Miljkovic-Lolic, M., Vanderhoek, J.Y., Fiskum, G., 1998. Normoxic ventilation after cardiac arrest reduces oxidation of brain lipids and improves neurological outcome. *Stroke* 29, 1679–1686.
- Lolic, M.M., Fiskum, G., Rosenthal, R.E., 1997. Neuroprotective effects of acetyl-L-carnitine after stroke in rats. *Ann. Emerg. Med.* 29, 758–765.
- MacDonald, H.M., Mulligan, J.C., Allen, A.C., Taylor, P.M., 1980. Neonatal asphyxia. I. Relationship of obstetric and neonatal complications to neonatal mortality in 38,405 consecutive deliveries. *J. Pediatr.* 96 (5), 898–902.
- Macri, M.A., Campanella, R., De, L.F., Montalbano, A., Taggi, F., Maraviglia, B., 1995. In vivo 31P spectroscopy study of treated and untreated recovery of rat partial brain ischemia. *Magn. Reson. Med.* 34, 542–547.
- Manole, M.D., Hickey, R.W., Clark, R.S., Kochanek, P.M., 2008. Current and future therapies of pediatric cardiopulmonary arrest. *Indian. J. Pediatr.* 75, 609–614.
- Martin, S.S., Perez-Polo, J.R., Noppens, K.M., Grafe, M.R., 2005. Biphasic changes in the levels of poly(ADP-ribose) polymerase-1 and caspase 3 in the immature brain following hypoxia-ischemia. *Int. J. Dev. Neurosci.* 23, 673–686.
- Mavelli, I., Rigo, A., Federico, R., Cirioli, M.R., Rotilio, G., 1982. Superoxide dismutase, glutathione peroxidase and catalase in developing rat brain. *Biochem. J.* 204, 535–540.
- Max, J.E., Koele, S.L., Castillo, C.C., Lindgren, S.D., Arndt, S., Bokura, H., Robin, D.A., Smith Jr., W.L., Sato, Y., 2000. Personality change disorder in children and adolescents following traumatic brain injury. *J. Int. Neuropsychol. Soc.* 6, 279–289.
- Max, J.E., Lansing, A.E., Koele, S.L., Bokura, H., Schachar, R., Collings, N., Williams, K.E., 2004. Attention deficit hyperactivity disorder in children and adolescents following traumatic brain injury. *Dev. Neuropsychol.* 25, 159–177.
- McDonald, J.W., Silverstein, F.S., Johnston, M.V., 1988. Neurotoxicity of N-methyl-D-aspartate is markedly enhanced in developing rat central nervous system. *Brain Res.* 459, 200–203.
- McDonald, J.W., Trescher, W.H., Johnston, M.V., 1992. Susceptibility of brain to AMPA induced excitotoxicity transiently peaks during early postnatal development. *Brain Res.* 583, 54–70.

- Milstein, J.M., White, J.G., Swaiman, K.F., 1968. Oxidative phosphorylation in mitochondria of developing rat brain. *J. Neurochem.* 15, 411–415.
- Monyer, H., Burnashev, N., Laurie, D.J., Sakmann, B., Seeburg, P.H., 1994. Developmental and regional expression in the rat brain and functional properties of four NMDA receptors. *Neuron* 12, 529–540.
- Murthy, M., Rappoport, D., 1963. Biochemistry of the developing rat brain. II. Neonatal mitochondrial oxidations. *Biochim. Biophys. Acta* 74, 51–59.
- Nakai, A., Shibasaki, Y., Taniuchi, Y., Miyake, H., Oya, A., Takeshita, T., 2004. Role of mitochondrial permeability transition in fetal brain damage in rats. *Pediatr. Neurol.* 30, 247–253.
- Nakajima, W., Ishida, A., Lange, M.S., Gabrielson, K.L., Wilson, M.A., Martin, L.J., Blue, M.E., Johnston, M.V., 2000. Apoptosis has a prolonged role in the neurodegeneration after hypoxic ischemia in the newborn rat. *J. Neurosci.* 20, 7994–8004.
- Natale, J.E., Joseph, J.G., Pretzlaff, R.K., Silber, T.J., Guerguerian, A.M., 2006. Clinical trials in pediatric traumatic brain injury: unique challenges and potential responses. *Dev. Neurosci.* 28, 276–290.
- Northington, F.J., Graham, E.M., Martin, L.J., 2005. Apoptosis in perinatal hypoxic-ischemic brain injury: how important is it and should it be inhibited? *Brain Res. Brain Res. Rev.* 50, 244–257.
- Perier, C., Tieu, K., Guegan, C., Caspersen, C., Jackson-Lewis, V., Carelli, V., Martinuzzi, A., Hirano, M., Przedborski, S., Vila, M., 2005. Complex I deficiency primes Bax-dependent neuronal apoptosis through mitochondrial oxidative damage. *Proc. Natl. Acad. Sci. U.S.A.* 102 (52), 19126–19131.
- Pohl, D., Bittigau, P., Ishimaru, M.J., Stadthaus, D., Hubner, C., Olney, J.W., Turski, L., Ikonomidou, C., 1999. N-Methyl-D-aspartate antagonists and apoptotic cell death triggered by head trauma in developing rat brain. *Proc. Natl. Acad. Sci. U.S.A.* 96, 2508–2513.
- Polster, B.M., Basanez, G., Etxebarria, A., Hardwick, J.M., Nicholls, D.G., 2005. Calpain I induces cleavage and release of apoptosis-inducing factor from isolated mitochondria. *J. Biol. Chem.* 280, 6447–6454.
- Polster, B.M., Basanez, G., Young, M., Suzuki, M., Fiskum, G., 2003a. Inhibition of Bax-induced cytochrome c release from neural cell and brain mitochondria by dibucaine and propranolol. *J. Neurosci.* 23, 2735–2743.
- Polster, B.M., Kinnally, K.W., Fiskum, G., 2001. Bcl-2 death domain peptide induces cell type-selective mitochondrial outer membrane permeability. *J. Biol. Chem.* 276 (41), 37887–37894.
- Polster, B.M., Robertson, C.L., Bucci, C.J., Suzuki, M., Fiskum, G., 2003c. Postnatal brain development and neural cell differentiation modulate mitochondrial Bax and BH3 peptide-induced cytochrome c release. *Cell Death Differ.* 10, 365–370.
- Polster, B.M., Robertson, C.L., Bucci, C.J., Suzuki, M., Fiskum, G., 2003b. Postnatal brain development and neural cell differentiation modulate mitochondrial Bax and BH3 peptide-induced cytochrome c release. *Cell Death Differ.* 10, 365–370.
- Prins, M.L., Fujima, L.S., Hovda, D.A., 2005. Age-dependent reduction of cortical contusion volume by ketones after traumatic brain injury. *J. Neurosci. Res.* 82, 413–420.
- Prins, M.L., Hovda, D.A., 2003. Developing experimental models to address traumatic brain injury in children. *J. Neurotrauma.* 20, 123–137.
- Puchowicz, M.A., Zechel, J.L., Valerio, J., Emancipator, D.S., Xu, K., Pundik, S., LaManna, J.C., Lust, W.D., 2008. Neuroprotection in diet-induced ketotic rat brain after focal ischemia. *J. Cereb. Blood Flow Metab.* 28, 1907–1916.
- Puka-Sundvall, M., Gajkowska, B., Cholewinski, M., Blomgren, K., Lazarewicz, J.W., Hagberg, H., 2000a. Subcellular distribution of calcium and ultrastructural changes after cerebral hypoxia-ischemia in immature rats. *Brain Res. Dev. Brain Res.* 125 (1–2), 31–41.
- Puka-Sundvall, M., Wallin, C., Gilland, E., Hallin, U., Wang, X., Sandberg, M., Karlsson, J., Blomgren, K., Hagberg, H., 2000b. Impairment of mitochondrial respiration after cerebral hypoxia-ischemia in immature rats: relationship to activation of caspase-3 and neuronal injury. *Brain Res. Dev. Brain Res.* 125, 43–50.
- Puka-Sundvall, M., Gilland, E., Hagberg, H., 2001. Cerebral hypoxia-ischemia in immature rats: involvement of mitochondrial permeability transition? *Dev. Neurosci.* 23, 192–197.
- Puka-Sundvall, M., Wallin, C., Gilland, E., Hallin, U., Wang, X., Sandberg, M., Karlsson, J., Blomgren, K., Hagberg, H., 2000c. Impairment of mitochondrial respiration after cerebral hypoxia-ischemia in immature rats: relationship to activation of caspase-3 and neuronal injury. *Brain Res. Dev. Brain Res.* 125, 43–50.
- Puka-Sundvall, M., Wallin, C., Gilland, E., Hallin, U., Wang, X., Sandberg, M., Karlsson, J., Blomgren, K., Hagberg, H., 2000d. Impairment of mitochondrial respiration after cerebral hypoxia-ischemia in immature rats: relationship to activation of caspase-3 and neuronal injury. *Brain Res. Dev. Brain Res.* 125 (1–2), 43–50.
- Rabi, Y., Rabi, D., Yee, W., 2007. Room air resuscitation of the depressed newborn: a systematic review and meta-analysis. *Resuscitation* 72, 353–363.
- Raff, M.C., Barres, B.A., Burne, J.F., Coles, H.S., Ishizaki, Y., Jacobson, M.D., 1993. Programmed cell death and the control of cell survival: lessons from the nervous system. *Science* 262, 695–700.
- Rajapakse, N., Shimizu, K., Kis, B., Snipes, J., Lacza, Z., Busija, D., 2002. Activation of mitochondrial ATP-sensitive potassium channels prevents neuronal cell death after ischemia in neonatal rats. *Neurosci. Lett.* 327, 208–212.
- Richards, E.M., Fiskum, G., Rosenthal, R.E., Hopkins, I., McKenna, M.C., 2007. Hyperoxic reperfusion after global ischemia decreases hippocampal energy metabolism. *Stroke* 38, 1578–1584.
- Richards, E.M., Rosenthal, R.E., Kristian, T., Fiskum, G., 2006. Postischemic hyperoxia reduces hippocampal pyruvate dehydrogenase activity. *Free Radic. Biol. Med.* 40, 1960–1970.
- Rizzuto, R., Simpson, A.W., Brini, M., Pozzan, T., 1992. Rapid changes of mitochondrial Ca<sup>2+</sup> revealed by specifically targeted recombinant aequorin [published erratum appears in *Nature* 1992 Dec 24–31;360(6406):768]. *Nature* 358 (6384), 325–327.
- Robertson, C.L., Bucci, C.J., Fiskum, G., 2004. Mitochondrial response to calcium in the developing brain. *Brain Res. Dev. Brain Res.* 151, 141–148.
- Robertson, C.M., Finer, N.N., Grace, M.G., 1989. School performance of survivors of neonatal encephalopathy associated with birth asphyxia at term. *J. Pediatr.* 114 (5), 753–760.
- Robertson, C.L., Saraswati, M., Fiskum, G., 2007. Mitochondrial dysfunction early after traumatic brain injury in immature rats. *J. Neurochem.* 101, 1248–1257.
- Robertson, C.L., Soane, L., Siegel, Z.T., Fiskum, G., 2006. The potential role of mitochondria in pediatric traumatic brain injury. *Dev. Neurosci.* 28, 432–446.
- Rosenthal, R.E., Fiskum, G., 1990. Brain mitochondrial function in cerebral ischemia and resuscitation. In: Schurr, A. (Ed.), *Cerebral Ischemia and Resuscitation*. CRC Press, New York, pp. 289–300.
- Rosenthal, R.E., Hamud, F., Fiskum, G., Varghese, P.J., Sharpe, S., 1987. Cerebral ischemia and reperfusion: prevention of brain mitochondrial injury by lidoflazine. *J. Cereb. Blood Flow Metab.* 7 (6), 752–758.
- Rosenthal, R.E., Williams, R., Bogaert, Y.E., Getson, P.R., Fiskum, G., 1992. Prevention of postischemic canine neurological injury through potentiation of brain energy metabolism by acetyl-L-carnitine. *Stroke* 23, 1312–1317.
- Sadanaga-Akiyoshi, F., Yao, H., Tanuma, S., Nakahara, T., Hong, J.S., Ibayashi, S., Uchimura, H., Fujishima, M., 2003. Nicotinamide attenuates focal ischemic brain injury in rats: with special reference to changes in nicotinamide and NAD<sup>+</sup> levels in ischemic core and penumbra. *Neurochem. Res.* 28, 1227–1234.
- Scaffidi, S., O'Brien, J., Hopkins, I., Robertson, C., Fiskum, G., McKenna, M., (2009) Delayed cerebral oxidative glucose metabolism after traumatic brain injury in young rats.
- Schinzl, A.C., Takeuchi, O., Huang, Z., Fisher, J.K., Zhou, Z., Rubens, J., Hetz, C., Danial, N.N., Moskowitz, M.A., Korsmeyer, S.J., 2005. Cyclophilin D is a component of mitochondrial permeability transition and mediates neuronal cell death after focal cerebral ischemia. *Proc. Natl. Acad. Sci. U.S.A.* 102 (34), 12005–12010.
- Setkowicz, Z., Ciarach, M., Guzik, R., Janeczko, K., 2004. Different effects of neuroprotectants FK-506 and cyclosporin A on susceptibility to pilocarpine-induced seizures in rats with brain injured at different developmental stages. *Epilepsy Res.* 61, 63–72.
- Shankaran, S., Laptok, A.R., Ehrenkranz, R.A., Tyson, J.E., McDonald, S.A., Donovan, E.F., Fanaroff, A.A., Poole, W.K., Wright, L.L., Higgins, R.D., Finer, N.N., Carlo, W.A., Duara, S., Oh, W., Cotten, C.M., Stevenson, D.K., Stoll, B.J., Lemons, J.A., Guillet, R., Jobe, A.H., 2005. Whole-body hypothermia for neonates with hypoxic-ischemic encephalopathy. *N. Engl. J. Med.* 353, 1574–1584.
- Sharma, R., Derr-Yellin, E.C., House, D.E., Kodavanti, P.R., 2000. Age-dependent effects of Aroclor 1254R on calcium uptake by subcellular organelles in selected brain regions of rats. *Toxicology* 156, 13–25.
- Shimohama, S., Fujimoto, S., Sumida, Y., Tanino, H., 1998. Differential expression of rat brain bcl-2 family proteins in development and aging. *Biochem. Biophys. Res. Commun.* 252, 92–96.
- Shuaib, A., Waqar, T., Wishart, T., Kanthan, R., Howlett, W., 1995. Acetyl-L-carnitine attenuates neuronal damage in gerbils with transient forebrain ischemia only when given before the insult. *Neurochem. Res.* 20, 1021–1025.
- Sims, N.R., 1991. Selective impairment of respiration in mitochondria isolated from brain subregions following transient forebrain ischemia in the rat. *J. Neurochem.* 56 (6), 1836–1844.
- Sitkiewicz, D., Skonieczna, M., Rychla, T., Gazdzik, D., Podymniak, S., Bicz, W., 1982. Modification of brain mitochondrial enzymes by the oxygen analogue of ronnel at various stages of development and aging. *J. Neurochem.* 39, 1308–1313.
- Soane, L., Siegel, Z.T., Schuh, R.A., Fiskum, G., 2008. Postnatal developmental regulation of Bcl-2 family proteins in brain mitochondria. *J. Neurosci. Res.* 86, 1267–1276.
- Stacpoole, P.W., 1989. The pharmacology of dichloroacetate. *Metabolism* 38, 1124–1144.
- Starkov, A.A., Chinopoulos, C., Fiskum, G., 2004. Mitochondrial calcium and oxidative stress as mediators of ischemic brain injury. *Cell Calcium* 36 (3–4), 257–264.
- Sullivan, P.G., Keller, J.N., Bussen, W.L., Scheff, S.W., 2002. Cytochrome c release and caspase activation after traumatic brain injury. *Brain Res.* 949 (1–2), 88–96.
- Sullivan, P.G., Rabchevsky, A.G., Waldmeier, P.C., Springer, J.E., 2005. Mitochondrial permeability transition in CNS trauma: cause or effect of neuronal cell death? *J. Neurosci. Res.* 79, 231–239.
- Suzuki, K., (1990) The changes of regional cerebral blood flow with advancing age in normal children. 159–170.
- Tai, K.K., Nguyen, N., Pham, L., Truong, D.D., 2008. Ketogenic diet prevents cardiac arrest-induced cerebral ischemic neurodegeneration. *J. Neural. Transm.* 115, 1011–1017.
- Tastekin, A., Gepdiremen, A., Ors, R., Emin, B.M., Halici, Z., 2005. L-Carnitine protects against glutamate- and kainic acid-induced neurotoxicity in cerebellar granular cell culture of rats. *Brain Res. Dev.* 27, 570–573.
- Taylor, A., Butt, W., Rosenfeld, J., Shann, F., Ditchfield, M., Lewis, E., Klug, G., Wallace, D., Henning, R., Tibballs, J., 2001. A randomized trial of very early decompressive craniectomy in children with traumatic brain injury and sustained intracranial hypertension. *Childs Nerv. Syst.* 17, 154–162.
- Thomas, S., Prins, M.L., Samii, M., Hovda, D.A., 2000. Cerebral metabolic response to traumatic brain injury sustained early in development: a 2-deoxy-D-glucose autoradiographic study. *J. Neurotrauma.* 17, 649–665.
- Tilford, J.M., Aitken, M.E., Anand, K.J., Green, J.W., Goodman, A.C., Parker, J.G., Killingsworth, J.B., Fiser, D.H., Adelson, P.D., 2005. Hospitalizations for critically ill children with traumatic brain injuries: a longitudinal analysis. *Crit. Care Med.* 33, 2074–2081.
- Tisdall, M.M., Tachtsidis, I., Leung, T.S., Elwell, C.E., Smith, M., 2008. Increase in cerebral aerobic metabolism by normobaric hyperoxia after traumatic brain injury. *J. Neurosurg.* 109, 424–432.
- Tolias, C.M., Reinert, M., Seiler, R., Gilman, C., Scharf, A., Bullock, M.R., 2004. Normobaric hyperoxia-induced improvement in cerebral metabolism and reduction in intracranial pressure in patients with severe head injury: a prospective historical cohort-matched study. *J. Neurosurg.* 101, 435–444.
- Tyler, D., Harrevel, A., 1942. The respiration of the developing brain. *Am. J. Physiol.* 136, 600–603.
- Uchino, H., Minamikawa-Tachino, R., Kristian, T., Perkins, G., Narazaki, M., Siesjo, B.K., Shibasaki, F., 2002. Differential neuroprotection by cyclosporin A and FK506

- following ischemia corresponds with differing abilities to inhibit calcineurin and the mitochondrial permeability transition. *Neurobiol. Dis.* 10 (3), 219–233.
- Vannucci, R.C., 1990. Current and potentially new management strategies for perinatal hypoxic-ischemic encephalopathy. *Pediatrics* 85 (6), 961–968.
- Vannucci, R.C., Duffy, T.E., 1976. Carbohydrate metabolism in fetal and neonatal rat brain during anoxia and recovery. *Am. J. Physiol.* 230, 1269–1275.
- Vannucci, S.J., Hagberg, H., 2004. Hypoxia–ischemia in the immature brain. *J. Exp. Biol.* 207, 3149–3154.
- Vannucci, S.J., Simpson, I.A., 2003. Developmental switch in brain nutrient transporter expression in the rat. *Am. J. Physiol. Endocrinol. Metab.* 285, E1127–E1134.
- Vannucci, R.C., Connor, J.R., Mauger, D.T., Palmer, C., Smith, M.B., Towfighi, J., Vannucci, S.J., 1999. Rat model of perinatal hypoxic-ischemic brain damage. *J. Neurosci. Res.* 55 (2), 158–163.
- Vesce, S., Jekabsons, M.B., Johnson-Cadwell, L.I., Nicholls, D.G., 2005. Acute glutathione depletion restricts mitochondrial ATP export in cerebellar granule neurons. *J. Biol. Chem.* 280 (46), 38720–38728.
- Vereczki, V., Martin, E., Rosenthal, R.E., Hof, P.R., Hoffman, G.E., Fiskum, G., 2006. Normoxic resuscitation after cardiac arrest protects against hippocampal oxidative stress, metabolic dysfunction, and neuronal death. *J. Cereb. Blood Flow Metab.* 26, 821–835.
- Wainwright, M.S., Kohli, R., Whittington, P.F., Chace, D.H., 2006. Carnitine treatment inhibits increases in cerebral carnitine esters and glutamate detected by mass spectrometry after hypoxia–ischemia in newborn rats. *Stroke* 37, 524–530.
- Wainwright, M.S., Mannix, M.K., Brown, J., Stumpf, D.A., 2003. L-Carnitine reduces brain injury after hypoxia–ischemia in newborn rats. *Pediatr. Res.* 54, 688–695.
- Walson, K.H., Clark, R.B., Tang, M., McFarland, A., Alexander, H., Kochanek, P., 2008. Normoxic versus hyperoxic resuscitation in pediatric asphyxial cardiac arrest: effects on oxidative stress. [abstract]. *Crit. Care Med.* 35 (12), A5.
- Wang, Q., Tang, X.N., Yenari, M.A., 2007. The inflammatory response in stroke. *J. Neuroimmunol.* 184 (1–2), 53–68.
- Wechsler, B., Kim, H., Gallagher, P.R., DiScala, C., Stineman, M.G., 2005. Functional status after childhood traumatic brain injury. *J. Trauma* 58, 940–949.
- Winter, B.K., Fiskum, G., Gallo, L.L., 1995. Effects of L-carnitine on serum triglyceride and cytokine levels in rat models of cachexia and septic shock. *Br. J. Cancer* 72, 1173–1179.
- Wu, L., Shen, F., Lin, L., Zhang, X., Bruce, I.C., Xia, Q., 2006. The neuroprotection conferred by activating the mitochondrial ATP-sensitive K<sup>+</sup> channel is mediated by inhibiting the mitochondrial permeability transition pore. *Neurosci. Lett.* 402, 184–189.
- Wyatt, J.S., Gluckman, P.D., Liu, P.Y., Azzopardi, D., Ballard, R., Edwards, A.D., Ferriero, D.M., Polin, R.A., Robertson, C.M., Thoresen, M., Whitelaw, A., Gunn, A.J., 2007. Determinants of outcomes after head cooling for neonatal encephalopathy. *Pediatrics* 119, 912–921.
- Ying, W., 2008. NAD<sup>+</sup> and NADH in ischemic brain injury. *Front Biosci.* 13, 1141–1151.
- Yager, J.Y., Brucklacher, R.M., Vannucci, R.C., 1991a. Cerebral oxidative metabolism and redox state during hypoxia–ischemia and early recovery in immature rats. *Am. J. Physiol.* 261, H1102–H1108.
- Yager, J.Y., Brucklacher, R.M., Vannucci, R.C., 1991b. Cerebral oxidative metabolism and redox state during hypoxia–ischemia and early recovery in immature rats. *Am. J. Physiol.* 261, H1102–H1108.
- Yager, J.Y., Brucklacher, R.M., Vannucci, R.C., 1996. Paradoxical mitochondrial oxidation in perinatal hypoxic–ischemic brain damage. *Brain Res.* 712, 230–238.
- Yamada, K.A., Rensing, N., Thio, L.L., 2005. Ketogenic diet reduces hypoglycemia-induced neuronal death in young rats. *Neurosci. Lett.* 385, 210–214.
- Yeates, K.O., Swift, E., Taylor, H.G., Wade, S.L., Drotar, D., Stancin, T., Minich, N., 2004. Short- and long-term social outcomes following pediatric traumatic brain injury. *J. Int. Neuropsychol. Soc.* 10, 412–426.
- Ying, W., Wei, G., Wang, D., Wang, Q., Tang, X., Shi, J., Zhang, P., Lu, H., 2007. Intranasal administration with NAD<sup>+</sup> profoundly decreases brain injury in a rat model of transient focal ischemia. *Front Biosci.* 12, 2728–2734.
- Zanelli, S.A., Solenski, N.J., Rosenthal, R.E., Fiskum, G., 2005. Mechanisms of Ischemic Neuroprotection by acetyl-L-carnitine. *Ann. N.Y. Acad. Sci.* 1053, 153–161.
- Zhang, L., Nair, A., Krady, K., Corpe, C., Bonneau, R.H., Simpson, I.A., Vannucci, S.J., 2004. Estrogen stimulates microglia and brain recovery from hypoxia–ischemia in normoglycemic but not diabetic female mice. *J. Clin. Invest.* 113, 85–95.
- Zhao, Q., Stafstrom, C.E., Fu, D.D., Hu, Y., Holmes, G.L., 2004. Detrimental effects of the ketogenic diet on cognitive function in rats. *Pediatr. Res.* 55, 498–506.
- Zhu, C., Qiu, L., Wang, X., Hallin, U., Cande, C., Kroemer, G., Hagberg, H., Blomgren, K., 2003. Involvement of apoptosis-inducing factor in neuronal death after hypoxia–ischemia in the neonatal rat brain. *J. Neurochem.* 86, 306–317.
- Zhu, C., Wang, X., Xu, F., Bahr, B.A., Shibata, M., Uchiyama, Y., Hagberg, H., Blomgren, K., 2005. The influence of age on apoptotic and other mechanisms of cell death after cerebral hypoxia–ischemia. *Cell Death Differ.* 12, 162–176.
- Zhu, H., Jia, Z., Strobl, J.S., Ehrlich, M., Misra, H.P., Li, Y., 2008. Potent induction of total cellular and mitochondrial antioxidants and phase 2 enzymes by cruciferous sulforaphane in rat aortic smooth muscle cells: cytoprotection against oxidative and electrophilic stress. *Cardiovasc. Toxicol.* 8, 115–125.
- Zwienerberg, M., Muizelaar, J.P., 1999. Severe pediatric head injury: the role of hyperemia revisited. *J. Neurotrauma* 16, 937–943.