

Effects of FK506 and Cyclosporin A on Calcium Ionophore-Induced Mitochondrial Depolarization and Cytosolic Calcium in Astrocytes and Neurons

Sibel Kahraman, Linda L. Bambrick, and Gary Fiskum^{*}

Department of Anesthesiology and the Center for Shock, Trauma and Anesthesiology Research, University of Maryland School of Medicine, Baltimore, Maryland

This study tested the hypothesis that sensitivity to the Ca^{2+} -induced loss of mitochondrial membrane potential ($\Delta\Psi_m$) and the sensitivity of the loss of $\Delta\Psi$ to mitochondrial permeability transition pore (PTP) inhibitors are different for neurons and astrocytes. Primary cultures of rat cortical neurons and astrocytes were exposed to the Ca^{2+} ionophore 4-Br-A23187, and $\Delta\Psi_m$ was monitored with the fluorescent probe tetramethylrhodamine methyl ester (TMRM). Ca^{2+} ionophore caused a decline in $\Delta\Psi_m$ in both cell types that was partially inhibited by cyclosporin A (CsA) in astrocytes but not in neurons. Another PTP inhibitor, 2-aminoethoxy-diphenylborate, was ineffective at protecting against mitochondrial depolarization, but depolarization was inhibited by FK506, an immunosuppressant drug similar to CsA that does not inhibit the PTP. CsA and FK506 both significantly reduced the ionophore-induced rise in $[\text{Ca}^{2+}]_i$ in both neurons and astrocytes. We conclude that the protective effects of CsA and FK506 against Ca^{2+} ionophore-induced mitochondrial membrane depolarization in intact astrocytes is not due to PTP inhibition but possibly is a consequence of inhibiting the rise in $[\text{Ca}^{2+}]_i$. © 2011 Wiley-Liss, Inc.

Key words: permeability transition pore; immunophilin; TMRM; calcium homeostasis; mitochondria; 4-Br-A23187

Cell death induced by a variety of insults is often preceded by an abnormal increase in $[\text{Ca}^{2+}]_i$. During cellular Ca^{2+} overload, mitochondria accumulate Ca^{2+} , which under some circumstances triggers opening of the inner membrane permeability transition pore (PTP). Opening of this nonselective pore results in the loss of mitochondrial membrane potential ($\Delta\Psi_m$); loss of Ca^{2+} and other mitochondrial solutes, including metabolic cofactors of <1,500 Da; and osmotic swelling (Zoratti and Szabo, 1995; Crompton, 1999; Di Lisa et al., 2001; Petronilli et al., 2001). The PTP is activated by Ca^{2+} and oxidative stress and inhibited by Mg^{2+} , adenine nucleotides, and the immunosuppressant drug cyclosporin A (CsA; Fournier et al., 1987; Gunter and Pfeiffer,

1990; Halestrap et al., 1997a,b; Kristal and Dubinsky, 1997). Brain mitochondria display heterogeneity in their response to Ca^{2+} -induced MPT, with neuronal mitochondria being relatively resistant to inhibition by CsA compared with astrocyte mitochondria (Brustovetsky and Dubinsky, 2000; Kristian et al., 2002; Bambrick et al., 2006). Given the importance of mitochondrial dysfunction in neural cell death, the cellular basis for the heterogeneous responses of brain mitochondria to large Ca^{2+} loads could provide insight into the mechanisms responsible for selective cell vulnerability and cell-selective targets of neuroprotective drugs, e.g., CsA (Uchino et al., 1998; Yoshimoto and Siesjo, 1999; Li et al., 2000; Sullivan et al., 2011). Previous experiments performed with digitonin-permeabilized primary cultures of cerebellar granule neurons and cortical astrocytes demonstrated that mitochondrial Ca^{2+} uptake, along with resistance to Ca^{2+} -induced bioenergetic dysfunction, was potentiated by CsA in astrocytes but not neurons (Bambrick et al., 2006). The much greater complexity of intact cells compared with isolated mitochondria or permeabilized cells can influence mitochondrial activities and their response to stress. The goal of this study was to test the hypothesis that mitochondria within intact neurons and astrocytes display differential sensitivity to inhibition of Ca^{2+} -induced PTP opening by CsA, as evaluated by measuring the response of $\Delta\Psi_m$ following exposure of cells to a Ca^{2+} ionophore. Comparisons of the effects of PTP inhibitors on the ionophore-induced rise in $[\text{Ca}^{2+}]_i$ uncovered an additional effect of immunosuppressant

Contract grant sponsor: DAMD; Contract grant number: 17-03-1-0745 (to L.L.B.); Contract grant sponsor: NIH; Contract grant numbers: NS07375 (to S.K.), NS34152 (to G.F.), HD016596 (to G.F.).

^{*}Correspondence to: Dr. Gary Fiskum, Department of Anesthesiology, University of Maryland School of Medicine, 685 W. Baltimore St., MSTF 5.34, Baltimore, MD 21201. E-mail: gfishum@anes.umm.edu

Received 25 August 2010; Revised 9 May 2011; Accepted 10 May 2011

Published online 11 July 2011 in Wiley Online Library (wileyonlinelibrary.com). DOI: 10.1002/jnr.22709

drugs that influences the interpretation of their effects on mitochondria and the PTP.

MATERIALS AND METHODS

Cell Culture

Cerebral cortex astrocytes and neurons were isolated from 17 day in utero Sprague-Dawley rat fetuses. All animal procedures were carried out according to the National Institutes of Health and the University of Maryland, Baltimore, guidelines for the care and use of laboratory animals. Cortical neurons were grown on 25-mm coverslips for 10–17 days in vitro, at a density of ~50,000 cells/coverslip, in Neurobasal medium with glutamine, penicillin, streptomycin, and B27 supplement in 95% air/5% CO₂ at 37°C. Glial proliferation was prevented by adding cytosine arabinofuranoside (5 μM) 24 hr after plating. Immunocytochemical measurements of glial fibrillary acid protein (GFAP) confirmed that cultures contained <1% glia. Cortical astrocytes were cultured in DMEM/F12 (1/1) 10% FBS with penicillin and streptomycin in 95% air/5% CO₂ at 37°C at a density of 20,000 cells/coverslip. For astrocytes, the culture medium was changed every 3 days, and the cells were used at 14–17 days in vitro, within 48 hr after changing the medium. Cultures were >95% astrocytes by GFAP immunocytochemistry.

Fluorescence Microscopy

Imaging of intracellular $\Delta\Psi_m$. $\Delta\Psi_m$ was followed in intact cortical neurons and astrocytes via epifluorescence microscopy. Cells were loaded with 50 nM tetramethylrhodamine methyl ester perchlorate (TMRM; Molecular Probes, Eugene, OR) in DMSO (0.1% final concentration) for 20 min at 37°C. The coverslips were mounted in the chamber of a Nikon Eclipse TE2000-S inverted microscope (objectives: SFluor ×20, 0.75 N.A., and ×40, 1.2 N.A.) in recording solution containing 120 mM NaCl, 3.5 mM KCl, 20 mM HEPES, 1.3 mM CaCl₂, 5 mM NaHCO₃, 1.0 mM MgCl₂, 15 mM glucose, and 5 nM TMRM, pH 7.4. Experiments were performed at room temperature. Single-cell fluorescence of TMRM was imaged by excitation at 547 nm (polychrome IV; Till, Munich, Germany) and emission at 620 nm. Image sequences (50-msec exposure time, 2 × 2 binning, 10-sec time lapse) were acquired from an ORCA-ER CCD camera (Hamamatsu Photonics, Hamamatsu, Germany). Data acquisition and analysis were conducted in Metafluor 5.0 (Universal Imaging, West Chester, PA) imaging software.

Ca²⁺ influx into the cells was stimulated by the addition of the calcium ionophore 4-bromo-A23187 (5 μM). The mitochondrial uncoupler FCCP (5 μM) was applied at the end of experiments to indicate the level of fluorescence of TMRM that reflected maximal dissipation of $\Delta\Psi_m$. Data are presented as TMRM fluorescence at time $t = [F_t - F_{FCCP}/F_0] \times 100\%$.

Imaging of intracellular Ca²⁺. Cells were loaded with 2 μM Fura FF (Molecular Probes) at room temperature in culture medium. Neurons were loaded for 20 min and astrocytes were loaded for 40 min, followed by a 15-min dye deesterification period. Cells were illuminated alternately at 340 and 380 nm, and fluorescence was monitored at 510 nm with the same imaging system as used for TMRM. Ratios

were calculated after subtraction of a background image from each wavelength (340 and 380 nm). [Ca²⁺]_i levels were expressed as the ratio of emitted fluorescence on excitation at 340 and 380 nm. Cells were preincubated with treatment drugs: 2 μM CsA, 1 μM FK506, and 100 μM 2-aminoethoxydiphenyl borate (2-APB) for 30 min. Drugs were also present in all solutions during the entire experiment.

Statistical Analysis

All data are expressed as mean ± SEM of n cells. Data from individual cultures were pooled to calculate group means. Data presented represent four to eight coverslips from two to four independent cell preparations. Statistical significance was assessed by one-way ANOVA test followed by the Tukey test for multiple comparisons. For data that were not normally distributed, the Mann-Whitney test or Kruskal-Wallis nonparametric ANOVA with Dunn's post hoc test for multiple comparisons of unbalanced data was used. $P < 0.05$ was considered to be statistically significant.

Materials

FK-506 was from LC Laboratories (Woburn, MA). All cell culture reagents were from Gibco-BRL (Grand Island, NY). All other chemical reagents were from Sigma (St. Louis, MO).

RESULTS

Ca²⁺ Ionophore-Induced Mitochondrial Depolarization in Cortical Astrocytes and Neurons

Baseline TMRM fluorescence intensity in both astrocytes and neurons was stable, declining less than 10% over 30 min, and was not affected by the presence of PTP inhibitors. Exposure of cells to 5 μM 4-Br-A23187 resulted in a rapid decline in TMRM fluorescence, as shown in Figure 1. Fluorescence changes were measured at 2, 5, and 10 min after the addition of the ionophore and came to completion by 10 min. Figure 2 compares these changes for astrocytes and neurons at 10 min after exposure to 4-Br-A23187. With ionophore alone, the decline in fluorescence intensity was approximately 80% for astrocytes and 75% for neurons. Preincubation of cells with CsA for 30 min attenuated the decrease in the astrocyte TMRM signal but actually increased the fluorescence loss in neurons (Figs. 1, 2). In addition to being a well-characterized inhibitor of the mitochondrial PTP, CsA is also a potent inhibitor of specific protein phosphatases, e.g., calcineurin. The effect of FK506 on ionophore-induced mitochondrial depolarization was tested as a control, because it also inhibits these phosphatases but does not inhibit the PTP (Conern and Halestrap, 1994). Surprisingly, FK506 also protected against the loss of TMRM fluorescence in astrocytes. As with CsA, FK506 also slightly promoted the drop in TMRM signal in neurons. The positive affect of FK506 with astrocytes suggests that PTP inhibition is not necessary for the inhibition of mitochondrial depolarization by CsA, so another PTP inhibitor without known effects on protein phosphorylation was tested. 2-Aminoethoxydiphenyl borate (2-APB) was shown pre-




Fig. 1. Ca^{2+} -induced loss of TMRM fluorescence in astrocytes and neurons. Mitochondrial membrane potential ($\Delta\Psi_m$) was followed in intact cortical neurons and astrocytes by epifluorescence microscopy with the fluorophore TMRM. Images are representative of those obtained for rat cortical astrocytes and neurons before and 10 min after the addition of the Ca^{2+} ionophore 4-Br-A23187 (5 μM) in the absence or presence of 45-min pretreatment and continued exposure to the PTP inhibitor cyclosporin A (CsA; 2 μM).

viously to inhibit PTP opening in brain mitochondria under conditions in which CsA is ineffective (Chinopoulos et al., 2003). This agent displayed no significant effect on the fluorescent decline with either astrocytes or neurons (Fig. 2).

Effects of Ca^{2+} Ionophore, CsA, and FK506 on Astrocyte and Neuronal $[\text{Ca}^{2+}]_i$

Additional experiments were performed to determine how CsA could protect against loss of $\Delta\Psi_m$ in astrocytes by a PTP-independent mechanism. The low-affinity Ca^{2+} -sensitive dye Fura FF AM was used to evaluate the large increases in $[\text{Ca}^{2+}]_i$ induced by exposure of astrocytes and neurons to 4-Br-A23187. Cortical astrocytes were heterogeneous in their response to 4-Br-A23187, with at least two distinct populations evident: those exhibiting a high initial peak and plateau (high responders) and those presenting a low transient initial peak followed by a delayed slight increase in $[\text{Ca}^{2+}]_i$ (low responders; Fig. 3A). Preincubation of astrocytes with CsA totally abolished the high responder response (Fig. 3B). Incubation with FK506 also reduced the number of cells that responded to the ionophore with a high initial




Fig. 2. Changes in mitochondrial membrane potential ($\Delta\Psi_m$) following exposure of astrocytes and neurons to Ca^{2+} ionophore. Images such as those in Figure 1 were analyzed for TMRM fluorescence intensity within the cell bodies of astrocytes and neurons before and 10 min after the addition of the Ca^{2+} ionophore 4-Br-A23187 (5 μM) in the absence (control) or presence of a 45-min pretreatment and continued exposure to cyclosporin A (CsA; 2 μM), FK506 (1 μM), or 2-aminoethoxydiphenyl borate (100 μM). Data are expressed as percentage of initial fluorescence before addition of ionophore, with 0% defined as the minimal fluorescence obtained 2 min after the addition of the mitochondrial uncoupler FCCP (5 μM) at 10 min after addition of ionophore. * $P < 0.05$ compared with control cells.

$[\text{Ca}^{2+}]_i$ peak (Fig. 3C). Addition of the mitochondrial respiratory uncoupler FCCP at the end of the experiments in which the initial rise in $[\text{Ca}^{2+}]_i$ was suppressed by CsA or FK506 resulted in an abrupt increase in $[\text{Ca}^{2+}]_i$ (Fig. 3B,C).

Exposure of cortical neurons to the Ca^{2+} ionophore induced a uniform rise in $[\text{Ca}^{2+}]_i$ that was more rapid than that observed in astrocytes (Fig. 4A). The presence of either CsA or FK506 slightly inhibited the rate and extent to which $[\text{Ca}^{2+}]_i$ rose in response to ionophore (Fig. 4B,C) but not to the extent observed for astrocytes (Fig. 3). The addition of FCCP resulted in a relatively very small additional rise in $[\text{Ca}^{2+}]_i$ under all conditions. Although there was no significant difference in the initial Fura FF ratio for astrocytes and neurons, the ratios observed after addition of ionophore in the absence or presence of CsA or FK506 were significantly greater for neurons than for astrocytes (compare Figs. 3A and 4A; $P < 0.001$).

DISCUSSION

The original aim of this study was to determine whether PTP opening in response to elevated $[\text{Ca}^{2+}]_i$ is different for astrocytes and neurons and whether the sensitivity to pharmacological inhibition is different between these two major types of brain cells. Although CsA inhibited the drop in $\Delta\Psi_m$ induced by Ca^{2+} ionophore in astrocytes, inhibition was also observed with FK506,




Fig. 3. Ca^{2+} ionophore-induced rise in $[\text{Ca}^{2+}]_i$ in cortical astrocytes. Changes in Fura FF fluorescence ratios (340 nm/380 nm) induced by 5 μM 4-Br-A23187 were heterogeneous among individual cortical astrocytes (light gray lines, $n = 100$), with two distinct populations evident, those with high initial peak and plateau and those with low transient initial peak followed by delayed slight increase in $[\text{Ca}^{2+}]_i$ (A). In contrast, the $[\text{Ca}^{2+}]_i$ increase after 4-Br-A23187 was uniformly low in all cells preincubated with CsA ($n = 134$; B). Incubation with FK506 also reduced the number of cells with high initial $[\text{Ca}^{2+}]_i$ peak ($n = 148$; C). Mean $[\text{Ca}^{2+}]_i$ values (solid lines with SEM [dark gray] in A–C) of CsA- and FK506-incubated cells were significantly lower than those of control cells. Five micromolar FCCP application 10 min after exposure to 4-Br-A23187 induced a further increase in treated cells (B,C).




Fig. 4. Ca^{2+} ionophore-induced rise in $[\text{Ca}^{2+}]_i$ in cortical neurons. Changes in Fura FF fluorescence ratios (340 nm/380 nm) following exposure of cortical neurons to 5 μM 4-Br-A23187 and 5 μM FCCP. 4-Br-A23187 caused a large, rapid increase in $[\text{Ca}^{2+}]_i$ in both drug-treated (B,C) and untreated (A) cells ($n = 211$). D provides a direct comparison between the mean increases (solid lines shown in A–C), demonstrating that the elevation observed in control cells was significantly higher than that of cells preincubated with 2 μM CsA (B; $n = 286$) or with 1 μM FK506 (C; $n = 166$).

an immunophilin binding drug that does not inhibit the PTP (Fig. 2). Moreover, inhibition was not observed with 2-APB, a PTP inhibitor that is a nonimmunophilin-binding drug (Fig. 2). Measurements of intracellular Ca^{2+} in response to Ca^{2+} ionophore demonstrated that both CsA and FK506 substantially dampened the rise in cytosolic Ca^{2+} caused by 4-Br-A23187 in astrocytes, indicating that this effect most likely is responsible for their protection against mitochondrial depolarization. In contrast to the effects of CsA and FK506 on astrocytes, these agents slightly promoted rather than inhibited neuronal mitochondrial depolarization in response to Ca^{2+} ionophore. It is also possible that these agents affected the TMRM measurements by influencing the membrane potential of the plasma membrane rather than that of the mitochondria; however, such effects would necessarily be different for the two cell types. Similar to the effects of these drugs on astrocyte $[\text{Ca}^{2+}]_i$, they slightly reduced the ionophore-induced rise in $[\text{Ca}^{2+}]_i$ in neurons.

These results fail to provide evidence for involvement of a CsA-sensitive PTP in mitochondrial depolarization in both astrocytes and neurons, albeit for different reasons. In astrocytes, the CsA inhibition of depolarization most likely is due to dampening the rise in intracellular Ca^{2+} , insofar as both effects are also observed with FK506. It is possible, however, that despite their effects on intracellular Ca^{2+} , these agents both inhibit mitochondrial depolarization that is independent of PTP or of changes in Ca^{2+} . These results do not exclude the possibility of a CsA-sensitive PTP but do reveal an additional mechanism of action for both CsA and FK506 that may obscure the direct effect of CsA on mitochondria. In neurons, neither CsA or FK506 protects against depolarization even though they reduce the rise in $[\text{Ca}^{2+}]_i$. It is possible, however, that CsA protection against loss of mitochondrial membrane potential in neurons would be observed at lower concentration of Ca^{2+} ionophore, because both brain and liver mitochondria lose their sensitivity to protection against PTP opening by CsA at very high Ca^{2+} loads (Chinopoulos et al., 2003). Also, it is possible that at high levels of Ca^{2+} ionophore, the cycling of Ca^{2+} in and out of the mitochondrial membrane could predominate over flux through the PTP, thus obscuring inhibition by PTP inhibitors. Moreover, our experiments utilized only a rise in intracellular Ca^{2+} to induce mitochondrial membrane depolarization, and the sensitivity to CsA could be different if oxidative stress plus Ca^{2+} were used as PTP inducers.

The ability of both CsA and FK506 to lower the large rise in $[\text{Ca}^{2+}]_i$ elicited by ionophore could be very important for interpreting the cytoprotective effects of these drugs in paradigms known to be associated with pathologic elevations of $[\text{Ca}^{2+}]_i$. Thus neuroprotection by both CsA and FK506 has been reported in animal models of stroke and traumatic brain injury, in which large elevations of intracellular Ca^{2+} are known to occur in both astrocytes and neurons (Kuroda et al., 1999; Singleton et al., 2001; Alessandri et al., 2002; Friberg and

Wieloch, 2002; Sullivan et al., 2011). Inhibition of the PTP or regulation of protein phosphorylation-dependent apoptotic pathways is typically invoked as the explanation for cytoprotection by immunophilin-based drugs. Our results indicate that their effects on intracellular Ca^{2+} should also be considered.

There are several possible mechanisms for the ability of these drugs to inhibit a rise in intracellular Ca^{2+} . They may relieve inhibition of noncapacitative Ca^{2+} channels (Mignen et al., 2003). CsA has also been reported to stimulate uptake of Ca^{2+} into the endoplasmic reticulum through the endoplasmic reticulum Ca^{2+} -ATPase (SERCA; Smaili et al., 2001). Although direct entry of Ca^{2+} into cells via 4-Br-A23187 is primarily responsible for the rise in $[\text{Ca}^{2+}]_i$ in our experiments, additional mechanisms, including those that both add to and buffer the rise in Ca^{2+} , can affect both the rate and the extent of Ca^{2+} elevation. It is also possible that CsA and FK506 could lower the effective concentration of the Ca^{2+} ionophore by stimulating its active efflux through the multidrug resistance transporter (Morjani and Madoulet, 2010); however, this mechanism is unlikely because of the high concentration of ionophore that was used and its strong affinity for cell membranes.

REFERENCES

- Alessandri B, Rice AC, Levasseur J, DeFord M, Hamm RJ, Bullock MR. 2002. Cyclosporin A improves brain tissue oxygen consumption and learning/memory performance after lateral fluid percussion injury in rats. *J Neurotrauma* 19:829–841.
- Bambrick LL, Chandrasekaran K, Mehrabian Z, Wright C, Krueger BK, Fiskum G. 2006. Cyclosporin A increases mitochondrial calcium uptake capacity in cortical astrocytes but not cerebellar granule neurons. *J Bioenerg Biomembr* 38:43–47.
- Brustovetsky N, Dubinsky JM. 2000. Dual responses of CNS mitochondria to elevated calcium. *J Neurosci* 20:103–113.
- Chinopoulos C, Starkov AA, Fiskum G. 2003. Cyclosporin A-insensitive permeability transition in brain mitochondria: inhibition by 2-aminooxydiphenyl borate. *J Biol Chem* 278:27382–27389.
- Connern CP, Halestrap AP. 1994. Recruitment of mitochondrial cyclophilin to the mitochondrial inner membrane under conditions of oxidative stress that enhance the opening of a calcium-sensitive nonspecific channel. *Biochem J* 302:321–324.
- Crompton M. 1999. The mitochondrial permeability transition pore and its role in cell death. *Biochem J* 341:233–249.
- Di Lisa F, Menabo R, Canton M, Barile M, Bernardi P. 2001. Opening of the mitochondrial permeability transition pore causes depletion of mitochondrial and cytosolic NAD^+ and is a causative event in the death of myocytes in postischemic reperfusion of the heart. *J Biol Chem* 276:2571–2575.
- Fournier N, Ducet G, Crevat A. 1987. Action of cyclosporine on mitochondrial calcium fluxes. *J Bioenerg Biomembr* 19:297–303.
- Friberg H, Wieloch T. 2002. Mitochondrial permeability transition in acute neurodegeneration. *Biochimie* 84:241–250.
- Gunter TE, Pfeiffer DR. 1990. Mechanisms by which mitochondria transport calcium. *Am J Physiol* 258:C755–C786.
- Halestrap AP, Connern CP, Griffiths EJ, Kerr PM. 1997a. Cyclosporin A binding to mitochondrial cyclophilin inhibits the permeability transition pore and protects hearts from ischaemia/reperfusion injury. *Mol Cell Biochem* 174:167–172.
- Halestrap AP, Woodfield KY, Connern CP. 1997b. Oxidative stress, thiol reagents, and membrane potential modulate the mitochondrial

- permeability transition by affecting nucleotide binding to the adenine nucleotide translocase. *J Biol Chem* 272:3346–3354.
- Kristal BS, Dubinsky JM. 1997. Mitochondrial permeability transition in the central nervous system: induction by calcium cycling-dependent and -independent pathways. *J Neurochem* 69:524–538.
- Kristian T, Weatherby TM, Bates TE, Fiskum G. 2002. Heterogeneity of the calcium-induced permeability transition in isolated non-synaptic brain mitochondria. *J Neurochem* 83:1297–1308.
- Kuroda S, Janelidze S, Siesjo BK. 1999. The immunosuppressants cyclosporin A and FK506 equally ameliorate brain damage due to 30-min middle cerebral artery occlusion in hyperglycemic rats. *Brain Res* 835:148–153.
- Li PA, Kristian T, He QP, Siesjo BK. 2000. Cyclosporin A enhances survival, ameliorates brain damage, and prevents secondary mitochondrial dysfunction after a 30-minute period of transient cerebral ischemia. *Exp Neurol* 165:153–163.
- Mignen O, Thompson JL, Shuttleworth TJ. 2003. Calcineurin directs the reciprocal regulation of calcium entry pathways in nonexcitable cells. *J Biol Chem* 278:40088–40096.
- Morjani H, Madoulet C. 2010. Immunosuppressors as multidrug resistance reversal agents. *Methods Mol Biol* 596:433–446.
- Petronilli V, Penzo D, Scorrano L, Bernardi P, Di LF. 2001. The mitochondrial permeability transition, release of cytochrome c and cell death. Correlation with the duration of pore openings in situ. *J Biol Chem* 276:12030–12034.
- Singleton RH, Stone JR, Okonkwo DO, Pellicane AJ, Povlishock JT. 2001. The immunophilin ligand FK506 attenuates axonal injury in an impact-acceleration model of traumatic brain injury. *J Neurotrauma* 18:607–614.
- Smaili SS, Stellato KA, Burnett P, Thomas AP, Gaspers LD. 2001. Cyclosporin A inhibits inositol 1,4,5-trisphosphate-dependent Ca^{2+} signals by enhancing Ca^{2+} uptake into the endoplasmic reticulum and mitochondria. *J Biol Chem* 276:23329–23340.
- Sullivan PG, Sebastian AH, Hall ED. 2011. Therapeutic window analysis of the neuroprotective effects of cyclosporine a after traumatic brain injury. *J Neurotrauma* 28:311–318.
- Uchino H, Elmer E, Uchino K, Li PA, He QP, Smith ML, Siesjo BK. 1998. Amelioration by cyclosporin A of brain damage in transient fore-brain ischemia in the rat. *Brain Res* 812:216–226.
- Yoshimoto T, Siesjo BK. 1999. Posttreatment with the immunosuppressant cyclosporin A in transient focal ischemia. *Brain Res* 839:283–291.
- Zoratti M, Szabo I. 1995. The mitochondrial permeability transition. *Biochim Biophys Acta* 1241:139–176.