

## 2012

116. Aschebrook-Kilfoy, B., Heltshe, S. L., Nuckols, J. R., et al. (2012). Modeled nitrate levels in well water supplies and prevalence of abnormal thyroid conditions among the Old Order Amish in Pennsylvania. *Environ Health* 11: 6.
117. Asselbergs, F. W., Guo, Y., van Iperen, E. P., et al. (2012). Large-scale gene-centric meta-analysis across 32 studies identifies multiple lipid loci. *Am J Hum Genet* 91(5): 823-838.
118. Billings, L. K., Hsu, Y. H., Ackerman, R. J., et al. (2012). Impact of common variation in bone-related genes on type 2 diabetes and related traits. *Diabetes* 61(8): 2176-2186.
119. Chasman, D. I., Fuchsberger, C., Pattaro, C., et al. (2012). Integration of genome-wide association studies with biological knowledge identifies six novel genes related to kidney function. *Hum Mol Genet* 21(24): 5329-5343.
120. Dastani, Z., Hivert, M. F., Timpson, N., et al. (2012). Novel loci for adiponectin levels and their influence on type 2 diabetes and metabolic traits: a multi-ethnic meta-analysis of 45,891 individuals. *PLoS Genet* 8(3): e1002607.
121. Ellis, J. W., Chen, M. H., Foster, M. C., et al. (2012). Validated SNPs for eGFR and their associations with albuminuria. *Hum Mol Genet* 21(14): 3293-3298.
122. Estrada, K., Styrkarsdottir, U., Evangelou, E., et al. (2012). Genome-wide meta-analysis identifies 56 bone mineral density loci and reveals 14 loci associated with risk of fracture. *Nat Genet* 44(5): 491-501.
123. Fox, C. S., Liu, Y., White, C. C., et al. (2012). Genome-wide association for abdominal subcutaneous and visceral adipose reveals a novel locus for visceral fat in women. *PLoS Genet* 8(5): e1002695.
124. Hu, Y., Snitker, S., Ryan, K. A., et al. (2012). Serum alanine aminotransferase is correlated with hematocrit in healthy human subjects. *Scand J Clin Lab Invest* 72(3): 258-264.
125. Lewis, J. P. and Shuldiner, A. R. (2012). Paraoxonase 1 Q192R variant and clopidogrel efficacy: fact or fiction? *Circ Cardiovasc Genet* 5(2): 153-155.
126. Liu, C. T., Estrada, K., Yerges-Armstrong, L. M., et al. (2012). Assessment of gene-by-sex interaction effect on bone mineral density. *J Bone Miner Res* 27(10): 2051-2064.
127. Liu, J., Hoppman, N., O'Connell, J. R., et al. (2012). A functional haplotype in EIF2AK3, an ER stress sensor, is associated with lower bone mineral density. *J Bone Miner Res* 27(2): 331-341.
128. Manning, A. K., Hivert, M. F., Scott, R. A., et al. (2012). A genome-wide approach accounting for body mass index identifies genetic variants influencing fasting glycemic traits and insulin resistance. *Nat Genet* 44(6): 659-669.
129. McArdle, P. F., Whitcomb, B. W., Tanner, K., et al. (2012). Association between bilirubin and cardiovascular disease risk factors: using Mendelian randomization to assess causal inference. *BMC Cardiovasc Disord* 12: 16.
130. Mitchell, B. D., Lee, W. J., Tolea, M. I., et al. (2012). Living the good life? Mortality and hospital utilization patterns in the Old Order Amish. *PLoS One* 7(12): e51560.
131. Mitchell, G. F., Verwoert, G. C., Tarasov, K. V., et al. (2012). Common genetic variation in the 3'-BCL11B gene desert is associated with carotid-femoral pulse wave velocity and

- excess cardiovascular disease risk: The AortaGen Consortium. *Circ Cardiovasc Genet* 5(1): 81-90.
- 132. Murabito, J. M., White, C. C., Kavousi, M., et al. (2012). Association between chromosome 9p21 variants and the ankle-brachial index identified by a meta-analysis of 21 genome-wide association studies. *Circ Cardiovasc Genet* 5(1): 100-112.
  - 133. Palmer, N. D., McDonough, C. W., Hicks, P. J., et al. (2012). A genome-wide association search for type 2 diabetes genes in African Americans. *PLoS One* 7(1): e29202.
  - 134. Parsa, A., Brown, E., Weir, M. R., et al. (2012). Genotype-based changes in serum uric acid affect blood pressure. *Kidney Int* 81(5): 502-507.
  - 135. Pattaro, C., Kottgen, A., Teumer, A., et al. (2012). Genome-wide association and functional follow-up reveals new loci for kidney function. *PLoS Genet* 8(3): e1002584.
  - 136. Saxena, R., Elbers, C. C., Guo, Y., et al. (2012). Large-scale gene-centric meta-analysis across 39 studies identifies type 2 diabetes loci. *Am J Hum Genet* 90(3): 410-425.
  - 137. Scott, R. A., Lagou, V., Welch, R. P., et al. (2012). Large-scale association analyses identify new loci influencing glycemic traits and provide insight into the underlying biological pathways. *Nat Genet* 44(9): 991-1005.
  - 138. Shuldiner, A. R., Vesely, M. R. and Fisch, A. (2012). CYP2C19 genotype and cardiovascular events. *JAMA* 307(14): 1482; author reply 1484-1485.
  - 139. Stolk, L., Perry, J. R., Chasman, D. I., et al. (2012). Meta-analyses identify 13 loci associated with age at menopause and highlight DNA repair and immune pathways. *Nat Genet* 44(3): 260-268.
  - 140. Wassel, C. L., Lamina, C., Nambi, V., et al. (2012). Genetic determinants of the ankle-brachial index: a meta-analysis of a cardiovascular candidate gene 50K SNP panel in the candidate gene association resource (CARe) consortium. *Atherosclerosis* 222(1): 138-147.
  - 141. Wilmot, B., Voruganti, V. S., Chang, Y. P., et al. (2012). Heritability of serum sodium concentration: evidence for sex- and ethnic-specific effects. *Physiol Genomics* 44(3): 220-228.
  - 142. Yang, J., Loos, R. J., Powell, J. E., et al. (2012). FTO genotype is associated with phenotypic variability of body mass index. *Nature* 490(7419): 267-272.
  - 143. Zheng, H. F., Tobias, J. H., Duncan, E., et al. (2012). WNT16 influences bone mineral density, cortical bone thickness, bone strength, and osteoporotic fracture risk. *PLoS Genet* 8(7): e1002745.
  - 144. Zupancic, M. L., Cantarel, B. L., Liu, Z., et al. (2012). Analysis of the gut microbiota in the Old Order Amish and its relation to the metabolic syndrome. *PLoS One* 7(8): e43052.